

HC 17/00/1051



6300/392  
4




INVESTOR IN PEOPLE

XX

# **PRIORITY DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

I also certify that by virtue of an assignment registered under the Patents Act 1977, the application is now proceeding in the name as substituted.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

*Andrew*

Dated

20 October 2000

---

**This Page Blank (uspto)**

GB9924092.1

---

By virtue of a direction given under Section of the Patents Act 1977, the application is proceeding in the name of

ASTRAZENECA AB,  
Incorporated in Sweden,  
S-151 85 Sodertalje,  
Sweden

[ADP No. 07822448003]

GB9924092.1

4/7/00  
By virtue of a direction given under Section of the Patents Act 1977, the application is proceeding in the name of

ASTRAZENECA UK LIMITED  
Incorporated in the United Kingdom  
15 Stanhope Gate  
LONDON  
W1Y 6AJ  
United Kingdom

[ADP No. 07810294001]

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road

Newport

Gwent NP9 1RH

1. Your reference

PHM 99-153

2. Patent application number

(The Patent Office will fill in this part)

9924092.1

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Zeneca Limited  
15 Stanhope Gate  
LONDON W1Y 6LN  
Great Britain

Patents ADP number (if you know it)

6254007002

If the applicant is a corporate body, give the country/state of its incorporation

SECTION 99 (1)(a) APPLICATION FILED 9/3/00  
THE PATENT OFFICE  
13 OCT 1999  
RECEIVED BY POST

4. Title of the invention

PYRIMIDINE DERIVATIVES

5. Name of your agent (if you have one)

TAIT, Brian Steele

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Global Intellectual Property, Patents  
AstraZeneca PLC  
Mereside, Alderley Park  
Macclesfield, Cheshire SK10 4TG  
Great Britain

Patents ADP number (if you know it)

5684600002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or

# Patents Form 1/77

Do not count copies of the same document

Continuation sheets of this form

Description

64

Claim(s)

Abstract

Drawing(s)

10. If you are also filing any of the following,  
state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right  
to grant of a patent (Patents Form 7/77)

Request for preliminary examination  
and search (Patents Form 9/77)

Request for substantive examination  
(Patents Form 10/77)

Any other documents  
(please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

*N. J. Marshall*

Date 12-Oct-99

Zeneca Limited Authorised Signatory

12. Name and daytime telephone number of  
person to contact in the United Kingdom

Joanne M Marshall

01625 516485

## Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

## Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.

**PYRIMIDINE DERIVATIVES**

This invention concerns certain novel pyrimidine derivatives and their use as inhibitors of cytokine mediated disease. The invention also concerns processes for the manufacture of  
5 said novel pyrimidine derivatives, pharmaceutical compositions containing them and their use in therapeutic methods, for example by virtue of inhibition of cytokine mediated disease.

The pyrimidine derivatives disclosed in the present invention are inhibitors of the  
production of cytokines such as Tumour Necrosis Factor (hereinafter TNF), for example  
TNF $\alpha$ , and various members of the interleukin (hereinafter IL) family, for example IL-1,  
10 IL-6 and IL-8. Accordingly the compounds of the invention will be useful in the treatment of diseases or medical conditions in which excessive production of cytokines occurs, for example excessive production of TNF $\alpha$  or IL-1. It is known that cytokines are produced by a wide variety of cells such as monocytes and macrophages and that they give rise to a variety of physiological effects which are believed to be important in disease or medical conditions  
15 such as inflammation and immunoregulation. For example, TNF $\alpha$  and IL-1 have been implicated in the cell signalling cascade which is believed to contribute to the pathology of disease states such as inflammatory and allergic diseases and cytokine-induced toxicity. It is also known that, in certain cellular systems, TNF $\alpha$  production precedes and mediates the production of other cytokines such as IL-1.

20 Abnormal levels of cytokines have also been implicated in, for example, the production of physiologically-active eicosanoids such as the prostaglandins and leukotrienes, the stimulation of the release of proteolytic enzymes such as collagenase, the activation of the immune system, for example by stimulation of T-helper cells, the activation of osteoclast activity leading to the resorption of calcium, the stimulation of the release of proteoglycans  
25 from, for example, cartilage, the stimulation of cell proliferation and to angiogenesis.

Cytokines are also believed to be implicated in the production and development of disease states such as inflammatory and allergic diseases, for example inflammation of the joints (especially rheumatoid arthritis, osteoarthritis and gout), inflammation of the gastrointestinal tract (especially inflammatory bowel disease, ulcerative colitis, Crohn's  
30 disease and gastritis), skin disease (especially psoriasis, eczema and dermatitis) and respiratory disease (especially asthma, bronchitis, allergic rhinitis and adult respiratory distress syndrome), and in the production and development of various cardiovascular and

cerebrovascular disorders such as congestive heart disease, myocardial infarction, the formation of atherosclerotic plaques, hypertension, platelet aggregation, angina, stroke, reperfusion injury, vascular injury including restenosis and peripheral vascular disease, and, for example, various disorders of bone metabolism such as osteoporosis (including senile and postmenopausal osteoporosis), Paget's disease, bone metastases, hypercalcaemia, hyperparathyroidism, osteosclerosis, osteoporosis and periodontitis, and the abnormal changes in bone metabolism which may accompany rheumatoid arthritis and osteoarthritis. Excessive cytokine production has also been implicated in mediating certain complications of bacterial, fungal and/or viral infections such as endotoxic shock, septic shock and toxic shock syndrome and in mediating certain complications of CNS surgery or injury such as neurotrauma and ischaemic stroke. Excessive cytokine production has also been implicated in mediating or exacerbating the development of diseases involving cartilage or muscle resorption, pulmonary fibrosis, cirrhosis, renal fibrosis, the cachexia found in certain chronic diseases such as malignant disease and acquired immune deficiency syndrome (AIDS), tumour invasiveness and tumour metastasis and multiple sclerosis.

Evidence of the central role played by  $\text{TNF}\alpha$  in the cell signalling cascade which gives rise to rheumatoid arthritis is provided by the efficacy in clinical studies of antibodies of  $\text{TNF}\alpha$  (The Lancet, 1994, 344, 1125 and British Journal of Rheumatology, 1995, 34, 334).

Thus cytokines such as  $\text{TNF}\alpha$  and IL-1 are believed to be important mediators of a considerable range of diseases and medical conditions. Accordingly it is expected that inhibition of the production of and/or effects of these cytokines will be of benefit in the prophylaxis, control or treatment of such diseases and medical conditions.


Without wishing to imply that the compounds disclosed in the present invention possess pharmacological activity only by virtue of an effect on a single biological process, it is believed that the compounds inhibit the effects of cytokines by virtue of inhibition of the enzyme p38 kinase. p38 kinase, otherwise known as cytokine suppressive binding protein (hereinafter CSBP) and reactivating kinase (hereinafter RK), is a member of the mitogen-activated protein (hereinafter MAP) kinase family of enzymes which is known to be activated by physiological stress such as that induced by ionising radiation, cytotoxic agents, and toxins, for example endotoxins such as bacterial lipopolysaccharide, and by a variety of agents such as the cytokines, for example  $\text{TNF}\alpha$  and IL-1. It is known that p38 kinase phosphorylates certain intracellular proteins which are involved in the cascade of enzymatic steps which leads




to the biosynthesis and excretion of cytokines such as  $\text{TNF}\alpha$  and IL-1. Known inhibitors of p38 kinase have been reviewed by G J Hanson in Expert Opinions on Therapeutic Patents, 1997, 7, 729-733. p38 kinase is known to exist in isoforms identified as p38 $\alpha$  and p38 $\beta$ .

The compounds disclosed in the present invention are inhibitors of the production of  
5 cytokines such as TNF, in particular of  $\text{TNF}\alpha$ , and various interleukins, in particular IL-1.

According to one aspect of the present invention there is provided a pyrimidine derivative of the Formula I



wherein m is 0, 1, 2 or 3 and each  $\text{R}^1$  group, which may be the same or different, is  
10 selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, sulphonamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-  
15 (2-6C)alkanoylamino, N-(1-6C)alkylsulphonamoyl, N,N-di-[(1-6C)alkyl]sulphonamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein  $\text{X}^1$  is a direct bond or is selected from O, S, SO,  $\text{SO}_2$ ,  $\text{N}(\text{R}^4)$ , CO,  $\text{CH}(\text{OR}^4)$ ,  
20  $\text{CON}(\text{R}^4)$ ,  $\text{N}(\text{R}^4)\text{CO}$ ,  $\text{SO}_2\text{N}(\text{R}^4)$ ,  $\text{N}(\text{R}^4)\text{SO}_2$ ,  $\text{OC}(\text{R}^4)_2$ ,  $\text{SC}(\text{R}^4)_2$  and  $\text{N}(\text{R}^4)\text{C}(\text{R}^4)_2$ , wherein each  $\text{R}^4$  is hydrogen or (1-6C)alkyl, and  $\text{Q}^2$  is aryl-(1-6C)alkyl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, or  $(\text{R}^1)_m$  is (1-3C)alkylenedioxy,

and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a  $\text{R}^1$  substituent is optionally separated by the insertion of a group selected from O, S, SO,  $\text{SO}_2$ ,  
25  $\text{N}(\text{R}^5)$ , CO,  $\text{CH}(\text{OR}^5)$ ,  $\text{CON}(\text{R}^5)$ ,  $\text{N}(\text{R}^5)\text{CO}$ ,  $\text{SO}_2\text{N}(\text{R}^5)$  and  $\text{N}(\text{R}^5)\text{SO}_2$  wherein  $\text{R}^5$  is hydrogen or (1-6C)alkyl,

and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl,

- 5 (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl,

---

N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :

- 10 
$$-X^2-Q^3$$

wherein X<sup>2</sup> is a direct bond or is selected from O and N(R<sup>7</sup>), wherein R<sup>7</sup> is hydrogen or (1-6C)alkyl, and Q<sup>3</sup> is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, and any Q<sup>3</sup> group optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, hydroxy, amino,

- 15 (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2 oxo or thioxo substituents,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group one or more halogeno or (1-6C)alkyl substituents or a substituent

- 20 selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl,
- 25 (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino;

R<sup>3</sup> is hydrogen, halogeno or (1-6C)alkyl;


n is 0, 1 or 2 and each R<sup>2</sup> group, which may be the same or different, is selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, (1-6C)alkoxycarbonyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and

- 30 di-[(1-6C)alkyl]amino;

p is 0, 1, 2, 3 or 4; and

Q<sup>1</sup> is aryl or heteroaryl and Q<sup>1</sup> is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino or with a (1-3C)alkylenedioxy group, or from a group of the

10 formula :



wherein X<sup>3</sup> is a direct bond or is selected from O and N(R<sup>8</sup>), wherein R<sup>8</sup> is hydrogen or (1-6C)alkyl, and Q<sup>4</sup> is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, and any Q<sup>4</sup> group optionally bears 1 or 2 substituents, which may

15 be the same or different, selected from halogeno, trifluoromethyl, cyano, hydroxy, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any heterocyclyl group within a substituent on Q<sup>1</sup> optionally bears 1 or 2 oxo or thioxo substituents,

and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a

20 Q<sup>1</sup> substituent is optionally separated by the insertion of a group selected from O, S, SO, SO<sub>2</sub>, N(R<sup>9</sup>), CO, CH(OR<sup>9</sup>), CON(R<sup>9</sup>), N(R<sup>9</sup>)CO, SO<sub>2</sub>N(R<sup>9</sup>) and N(R<sup>9</sup>)SO<sub>2</sub> wherein R<sup>9</sup> is hydrogen or (1-6C)alkyl,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a Q<sup>1</sup> group optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group one or more halogeno or (1-6C)alkyl substituents or a substituent selected

25 from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino;

30 or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof.

In this specification, the term (1-6C)alkyl includes straight and branched chain alkyl groups such as propyl, isopropyl and tert-butyl, unsaturated alkyl groups, for example (2-6C)alkenyl groups such as vinyl and allyl and (2-6C)alkynyl groups such as ethynyl and propargyl, (3-6C)cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl and  
 5 cyclohexyl and (3-6C)cycloalkenyl groups such as cyclopentenyl and cyclohexenyl.

References to individual alkyl groups such as propyl are specific for the straight chain version only and references to individual branched chain alkyl groups such as isopropyl are specific

---

for the branched chain version only. A similar convention applies to other generic groups, for example (1-6C)alkoxy includes methoxy, ethoxy, vinyloxy, allyloxy, propargyloxy,

10 cyclopropyloxy and cyclopentyloxy, (1-6C)alkylamino includes methylamino, ethylamino, allylamino, propargylamino, cyclobutylamino and cyclohexylamino, di-[(1-6Calkyl)amino includes dimethylamino, diethylamino, N-allyl-N-methylamino, N-methyl-N-propargylamino, N-cyclobutyl-N-methylamino and N-cyclohexyl-N-ethylamino and (2-6C)alkanoylamino includes acetamido, propionamido, acrylamido and propiolamido.

15 It is to be understood that, insofar as certain of the compounds of Formula I defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the above-mentioned activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for  
 20 example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory techniques referred to hereinafter.

Suitable values for the generic radicals referred to above include those set out below.

A suitable value for any one of the 'Q' groups ( $Q^1$  to  $Q^4$ ) when it is aryl or for the aryl  
 25 group within a 'Q' group is, for example, phenyl or naphthyl, preferably phenyl.

A suitable value for any one of the 'Q' groups ( $Q^1$  to  $Q^4$ ) when it is heteroaryl or for the heteroaryl group within a 'Q' group is, for example, an aromatic 5- or 6-membered monocyclic ring, a 9- or 10-membered bicyclic ring or a 13- or 14-membered tricyclic ring each containing 1 oxygen heteroatom or 1 or 2 nitrogen heteroatoms and optionally containing  
 30 a further heteroatom selected from nitrogen, oxygen and sulphur, for example furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazenyl,

benzofuranyl, indolyl, benzothiophenyl, benzoxazolyl, benzimidazolyl, benzothiazolyl, indazolyl, benzofurazanyl, quinolyl, isoquinolyl, quinazolinyl, quinoxalinyl, cinnolinyl, naphthyridinyl, carbazolyl, dibenzofuranyl, dibenzothiophenyl, S,S-dioxodibenzothiophenyl, xanthenyl, dibenzo-1,4-dioxinyl, phenoxathiinyl, phenoxazinyl, dibenzothiinyl, 5 phenothiazinyl, thianthrenyl, benzofuropyridyl, pyridoindolyl, acridinyl or phenanthridinyl, preferably furyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, benzofuranyl, indolyl, benzothiophenyl, benzoxazolyl, benzimidazolyl, benzothiazolyl, indazolyl, benzofurazanyl, quinolyl, isoquinolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, carbazolyl, dibenzofuranyl, 10 dibenzothiophenyl or xanthenyl, more preferably furyl, thienyl, isoxazolyl, thiazolyl, pyridyl, benzothiophenyl, benzofurazanyl, quinolyl, carbazolyl, dibenzofuranyl or dibenzothiophenyl, even more preferably pyridyl.

A suitable value for any one of the 'Q' groups ( $Q^1$  to  $Q^4$ ) when it is heterocyclyl or for the heterocyclyl group within a 'Q' group is, for example, a non-aromatic saturated or 15 partially saturated 3- to 10-membered monocyclic or bicyclic ring with up to three heteroatoms selected from oxygen, nitrogen and sulphur, for example oxiranyl, oxetanyl, azetidiny, tetrahydrofuranyl, tetrahydropyranyl, pyrrolinyl, pyrrolidinyl, morpholinyl, tetrahydro-1,4-thiazinyl, 1,1-dioxotetrahydro-1,4-thiazinyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperazinyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl or 20 tetrahydropyrimidinyl, preferably azetidin-1-yl, 3-pyrrolin-1-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, morpholino, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yl, piperidin-4-yl, homopiperidin-1-yl, piperazin-1-yl or homopiperazin-1-yl. A suitable value for such a group which bears 1 or 2 oxo or thioxo substituents is, for example, 2-oxopyrrolidinyl, 2-thioxopyrrolidinyl, 2-oxoimidazolidinyl, 2-thioxoimidazolidinyl, 25 2-oxopiperidinyl, 2,5-dioxopyrrolidinyl, 2,5-dioxoimidazolidinyl or 2,6-dioxopiperidinyl.

A suitable value for a 'Q' group when it is heteroaryl-(1-6C)alkyl is, for example, heteroarylmethyl, 2-heteroarylethyl and 3-heteroarylpropyl. The invention comprises corresponding suitable values for 'Q' groups when, for example, rather than a heteroaryl-(1-6C)alkyl group, an aryl-(1-6C)alkyl or heterocyclyl-(1-6C)alkyl group is present.

30 Suitable values for any of the 'R' groups ( $R^1$  to  $R^9$ ), or for various groups within an  $R^1$  substituent, or within a substituent on  $Q^1$  include:-

for halogeno

fluoro, chloro, bromo and iodo;

	for (1-6C)alkyl:	methyl, ethyl, vinyl, ethynyl, propyl, allyl, propargyl, isopropyl, cyclopropyl, <u>tert</u> -butyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl and cyclohexenyl;
5	for (1-6C)alkoxy:	methoxy, ethoxy, vinyloxy, propoxy, allyloxy, isopropoxy, propargyloxy, cyclopropyloxy, butoxy and cyclopentyloxy;
	for (1-6C)alkylthio:	methylthio, ethylthio and propylthio;
	for (1-6C)alkylsulphinyl:	methylsulphinyl and ethylsulphinyl;
10	for (1-6C)alkylsulphonyl:	methylsulphonyl and ethylsulphonyl;
	for (1-6C)alkylamino:	methylamino, ethylamino, propylamino, allylamino, propargylamino, isopropylamino, butylamino, cyclobutylamino and cyclohexylamino;
15	for di-[(1-6C)alkyl]amino:	dimethylamino, diethylamino, <u>N</u> -ethyl- <u>N</u> -methylamino, diisopropylamino, <u>N</u> -allyl- <u>N</u> -methylamino, <u>N</u> -methyl- <u>N</u> -propargylamino, <u>N</u> -cyclobutyl- <u>N</u> -methylamino and <u>N</u> -cyclohexyl- <u>N</u> -ethylamino;
20	for (1-6C)alkoxycarbonyl:	methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl and <u>tert</u> -butoxycarbonyl;
	for <u>N</u> -(1-6C)alkylcarbamoyl:	<u>N</u> -methylcarbamoyl, <u>N</u> -ethylcarbamoyl and <u>N</u> -propylcarbamoyl;
	for <u>N,N</u> -di-[(1-6C)alkyl]carbamoyl:	<u>N,N</u> -dimethylcarbamoyl, <u>N</u> -ethyl- <u>N</u> -methylcarbamoyl and <u>N,N</u> -diethylcarbamoyl;
25	for (2-6C)alkanoyl:	acetyl and propionyl;
	for (2-6C)alkanoyloxy:	acetoxyl and propionyloxy;
	for (2-6C)alkanoylamino:	acetamido, propionamido, acrylamido, methacrylamido, crotonamido and propiolamido;
30	for <u>N</u> -(1-6C)alkyl-(2-6C)alkanoylamino:	<u>N</u> -methylacetamido, <u>N</u> -methylpropionamido, <u>N</u> -methylacrylamido and <u>N</u> -methylpropiolamido;
	for <u>N</u> -(1-6C)alkylsulphamoyl:	<u>N</u> -methylsulphamoyl and <u>N</u> -ethylsulphamoyl;
	for <u>N,N</u> -di-[(1-6C)alkyl]sulphamoyl:	<u>N,N</u> -dimethylsulphamoyl;

for (1-6C)alkanesulphonylamino: methanesulphonylamino, ethanesulphonylamino,  
vinylsulphonylamino and ethynylsulphonylamino;

for N-(1-6C)alkyl-(1-6C)alkanesulphonylamino: N-methylmethanesulphonylamino,  
N-methylethanesulphonylamino,  
5 N-methylvinylsulphonylamino and  
N-methylethynylsulphonylamino.

A suitable value for  $(R^1)_m$  or for a substituent on  $Q^1$  when it is (1-3C)alkylenedioxy is,  
for example, methylenedioxy or ethylenedioxy and the oxygen atoms thereof occupy adjacent  
ring positions.

10 When, as defined hereinbefore, an  $R^1$  group forms a group of the formula  $Q^2-X^1$ - and,  
for example,  $X^1$  is a  $OC(R^4)_2$  linking group, it is the carbon atom, not the oxygen atom, of the  
 $OC(R^4)_2$  linking group which is attached to the pyrimidine ring of Formula I and the oxygen  
atom is attached to the  $Q^2$  group. A similar convention applies to the attachment of other  
groups of the formula  $Q^2-X^1$ -.

15 As defined hereinbefore, a single pair of adjacent carbon atoms in a (2-6C)alkylene  
chain within a  $R^1$  substituent is optionally separated by the insertion of a group such as O or  
 $CON(R^5)$  and a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a  $Q^1$   
substituent is optionally separated by the insertion of a group such as O or  $CON(R^9)$ . For  
example, insertion of a CONH group into the propylene chain within a 3-methoxypropoxy  
20 group gives rise to, for example, a 2-(2-methoxyacetamido)ethoxy group.

When, as defined hereinbefore, any  $CH_2$  or  $CH_3$  group within a  $R^1$  substituent  
optionally bears on each said  $CH_2$  or  $CH_3$  group one or more halogeno substituents, there are  
suitably 1 or 2 halogeno substituents present on each said  $CH_2$  group and there are suitably  
1, 2 or 3 halogeno substituents present on each said  $CH_3$  group.

25 When, as defined hereinbefore, any  $CH_2$  or  $CH_3$  group within a  $R^1$  substituent  
optionally bears on each said  $CH_2$  or  $CH_3$  group a substituent as defined hereinbefore, suitable  
 $R^1$  substituents so formed include, for example, hydroxy-substituted heterocyclyl-  
(1-6C)alkoxy groups such as 2-hydroxy-3-piperidinopropoxy and 2-hydroxy-  
3-morpholinopropoxy, hydroxy-substituted amino-(2-6C)alkoxy groups such as 3-amino-  
30 2-hydroxypropoxy, hydroxy-substituted (1-6C)alkylamino-(2-6C)alkoxy groups such as  
2-hydroxy-3-methylaminopropoxy, hydroxy-substituted di-[(1-6C)alkyl]amino-(2-6C)alkoxy  
groups such as 3-dimethylamino-2-hydroxypropoxy, hydroxy-substituted heterocyclyl-

(1-6C)alkylamino groups such as 2-hydroxy-3-piperidinopropylamino and 2-hydroxy-3-morpholinopropylamino, hydroxy-substituted amino-(2-6C)alkylamino groups such as 3-amino-2-hydroxypropylamino, hydroxy-substituted (1-6C)alkylamino-(2-6C)alkylamino groups such as 2-hydroxy-3-methylaminopropylamino, hydroxy-substituted

5 di-[(1-6C)alkyl]amino-(2-6C)alkylamino groups such as 3-dimethylamino-2-hydroxypropylamino, hydroxy-substituted (1-6C)alkoxy groups such as 2-hydroxyethoxy, (1-6C)alkoxy-substituted (1-6C)alkoxy groups such as 2-methoxyethoxy and

---

3-ethoxypropoxy and (1-6C)alkylsulphonyl-substituted (1-6C)alkoxy groups such as 2-methylsulphonylethoxy.

10 A suitable pharmaceutically-acceptable salt of a compound of the Formula I is, for example, an acid-addition salt of a compound of the Formula I, for example an acid-addition salt with an inorganic or organic acid such as hydrochloric, hydrobromic, sulphuric, trifluoroacetic, citric or maleic acid; or, for example, a salt of a compound of the Formula I which is sufficiently acidic, for example an alkali or alkaline earth metal salt such as a

15 calcium or magnesium salt, or an ammonium salt, or a salt with an organic base such as methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Various forms of prodrugs are known in the art. For examples of such prodrug derivatives, see:

- 20 a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- 25 c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- d) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- e) N. Kakeya, *et al.*, Chem. Pharm. Bull., 32, 692 (1984).


Examples of such pro-drugs may be used to form in-vivo-cleavable esters of a compound of the Formula I. An in-vivo-cleavable ester of a compound of the Formula I

30 containing a carboxy group is, for example, a pharmaceutically-acceptable ester which is cleaved in the human or animal body to produce the parent acid. Suitable pharmaceutically-acceptable esters for carboxy include (1-6C)alkoxymethyl esters, for example



- methoxymethyl; (1-6C)alkanoyloxymethyl esters, for example pivaloyloxymethyl; phthalidyl esters; (3-8C)cycloalkoxycarbonyloxy(1-6C)alkyl esters, for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolan-2-ylmethyl esters, for example 5-methyl-1,3-dioxolan-2-ylmethyl; and (1-6C)alkoxycarbonyloxyethyl esters, for example 1-methoxycarbonyloxyethyl; and may be formed at any carboxy group in the compounds of this invention.

- Particular novel compounds of the invention include, for example, pyrimidine derivatives of the Formula I, or pharmaceutically-acceptable salts thereof, wherein, unless otherwise stated, each of  $m$ ,  $R^1$ ,  $n$ ,  $R^2$ ,  $R^3$ ,  $p$  and  $Q^1$  has any of the meanings defined hereinbefore or in relevant paragraphs selected from paragraphs (a) to (u) hereinafter :-
- (a)  $m$  is 0, 1, 2 or 3, and each  $R^1$  group, which may be the same or different, is selected from hydroxy, halogeno, trifluoromethyl, amino, carbamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoylamino and N-(1-6C)alkyl-(2-6C)alkanoylamino,
- or from a group of the formula :



- wherein  $X^1$  is a direct bond or is selected from O,  $N(R^4)$ ,  $CON(R^4)$ ,  $N(R^4)CO$  and  $OC(R^4)_2$  wherein  $R^4$  is hydrogen or (1-6C)alkyl, and  $Q^2$  is aryl-(1-6C)alkyl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,
- and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a  $R^1$  substituent is optionally separated by the insertion of a group selected from O,  $N(R^5)$ ,  $CON(R^5)$  and  $N(R^5)CO$  wherein  $R^5$  is hydrogen or (1-6C)alkyl,
- and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on  $R^1$  optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, hydroxy, amino, carboxy, (1-6C)alkyl, (1-6C)alkoxy and (1-6C)alkoxycarbonyl,
- and wherein any heterocyclyl group within a substituent on  $R^1$  optionally bears 1 or 2 oxo substituents,
- and wherein any  $CH_2$  or  $CH_3$  group within a  $R^1$  substituent optionally bears on each said  $CH_2$  or  $CH_3$  group one or more (1-6C)alkyl substituents or a substituent selected from

hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

- (b) m is 0, 1, 2 or 3, and each R<sup>1</sup> group, which may be the same or different, is selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl, vinyl, ethynyl, methoxy, ethoxy, propoxy, methylthio, ethylthio, methylsulphinyl, methylsulphonyl, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, propargylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl, acetamido, propionamido, acrylamido and propiolamido, or from a group of the formula :



- wherein X<sup>1</sup> is a direct bond or is selected from O, NH, N(Me), CONH, NHCO and OCH<sub>2</sub> and Q<sup>2</sup> is benzyl, furylmethyl, thienylmethyl, imidazolymethyl, 2-imidazoleethyl, 3-imidazolepropyl, 4-imidazolebutyl, oxazolymethyl, thiazolymethyl, 1,2,3-triazolymethyl, 2-(1,2,3-triazolyl)ethyl, 3-(1,2,3-triazolyl)propyl, 1,2,4-triazolymethyl, 2-(1,2,4-triazolyl)ethyl, 3-(1,2,4-triazolyl)propyl, pyridylmethyl, 2-pyridylethyl, 3-pyridylpropyl, pyrrolidinyl, morpholinyl, tetrahydro-4H-1,4-thiazinyl, 1,1-dioxotetrahydro-4H-1,4-thiazinyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperazinyl, pyrrolidinylmethyl, 2-pyrrolidineethyl, 3-pyrrolidinylpropyl, imidazolidinylmethyl, 2-imidazolidinylethyl, 3-imidazolidinylpropyl, morpholinylmethyl, 2-morpholineethyl, 3-morpholinylpropyl, tetrahydro-4H-1,4-thiazinylmethyl, 1,1-dioxotetrahydro-4H-1,4-thiazinylmethyl, 2-(tetrahydro-4H-1,4-thiazinyl)ethyl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazinyl)ethyl, 3-(tetrahydro-4H-1,4-thiazinyl)propyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazinyl)propyl, piperidinylmethyl, 2-piperidineethyl, 3-piperidinylpropyl, homopiperidinylmethyl, 2-homopiperidineethyl, 3-homopiperidinylpropyl, piperazinylmethyl, 2-piperazineethyl, 3-piperazinylpropyl, homopiperazinylmethyl, 2-homopiperazineethyl or 3-homopiperazinylpropyl,

and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a R<sup>1</sup> substituent is optionally separated by the insertion of a group selected from O, NH, CONH and NHCO,

- 30 and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from

hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl, and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2 oxo substituents,

- 5 and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and diethylamino;

- (c) m is 0, 1, 2 or 3, and each R<sup>1</sup> group, which may be the same or different, is selected  
 10 from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylthio, methylsulphanyl, methylsulphonyl, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, propargylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl and acetamido, or from a group of the formula :

- 15 
$$Q^2 - X^1 -$$

- wherein X<sup>1</sup> is a direct bond or is selected from O, NH, N(Me), CONH, NHCO and OCH<sub>2</sub> and Q<sup>2</sup> is benzyl, 2-furylmethyl, 3-furylmethyl, 2-thienylmethyl, 3-thienylmethyl, 1-imidazolylmethyl, 2-imidazolylmethyl, 2-imidazol-1-ylethyl, 3-imidazol-1-ylpropyl, 4-imidazol-1-ylbutyl, 2-oxazolylmethyl, 4-oxazolylmethyl, 5-oxazolylmethyl,  
 20 2-thiazolylmethyl, 4-thiazolylmethyl, 5-thiazolylmethyl, 1,2,3-triazol-1-ylmethyl, 2-(1,2,3-triazol-1-yl)ethyl, 3-(1,2,3-triazol-1-yl)propyl, 1,2,4-triazol-1-ylmethyl, 2-(1,2,4-triazol-1-yl)ethyl, 3-(1,2,4-triazol-1-yl)propyl, 2-pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl, 2-pyrid-2-ylethyl, 2-pyrid-3-ylethyl, 2-pyrid-4-ylethyl, 3-pyrid-2-ylpropyl, 3-pyrid-3-ylpropyl, 3-pyrid-4-ylpropyl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl,  
 25 morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, homopiperidin-1-yl, homopiperidin-2-yl, homopiperidin-3-yl, homopiperidin-4-yl, piperazin-1-yl, homopiperazin-1-yl, pyrrolidin-1-ylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-1-ylpropyl, pyrrolidin-2-ylmethyl, 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-2-ylpropyl,  
 30 pyrrolidin-3-ylmethyl, 2-pyrrolidin-3-ylethyl, 3-pyrrolidin-3-ylpropyl, imidazolidin-1-ylmethyl, 2-imidazolidin-1-ylethyl, 3-imidazolidin-1-ylpropyl, imidazolidin-2-ylmethyl, 2-imidazolidin-2-ylethyl, 3-imidazolidin-2-ylpropyl,

- morpholinomethyl, 2-morpholinoethyl, 3-morpholinopropyl, morpholin-2-ylmethyl, 2-morpholin-2-ylethyl, 3-morpholin-2-ylpropyl, morpholin-3-ylmethyl, 2-morpholin-3-ylethyl, 3-morpholin-3-ylpropyl, tetrahydro-4H-1,4-thiazin-4-ylmethyl, 2-(tetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(tetrahydro-4H-1,4-thiazin-4-yl)propyl,
- 5 1,1-dioxotetrahydro-4H-1,4-thiazin-4-ylmethyl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propyl, piperidinomethyl, 2-piperidinoethyl, 3-piperidinopropyl, piperidin-2-ylmethyl, 2-piperidin-2-ylethyl, 3-piperidin-2-ylpropyl, piperidin-3-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-3-ylpropyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl,
- 10 homopiperidin-1-ylmethyl, 2-homopiperidin-1-ylethyl, 3-homopiperidin-1-ylpropyl, homopiperidin-2-ylmethyl, 2-homopiperidin-2-ylethyl, 3-homopiperidin-2-ylpropyl, homopiperidin-3-ylmethyl, 2-homopiperidin-3-ylethyl, 3-homopiperidin-3-ylpropyl, homopiperidin-4-ylmethyl, 2-homopiperidin-4-ylethyl, 3-homopiperidin-4-ylpropyl, piperazin-1-ylmethyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, piperazin-2-ylmethyl,
- 15 2-piperazin-2-ylethyl, 3-piperazin-2-ylpropyl, homopiperazin-1-ylmethyl, 2-homopiperazin-1-ylethyl, 3-homopiperazin-1-ylpropyl, homopiperazin-2-ylmethyl, 2-homopiperazin-2-ylethyl or 3-homopiperazin-2-ylpropyl,
- and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a R<sup>1</sup> substituent is optionally separated by the insertion of a group selected from O, NH, CONH
- 20 and NHCO,
- and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,
- 25 and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2 oxo substituents;
- and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and
- 30 diethylamino;
- (d) m is 0, 1 or 2 and each R<sup>1</sup> group, which may be the same or different, is selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl,

- methoxy, ethoxy, propoxy, methylthio, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, propargylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl, acetamido, benzyl, benzyloxy, benzylamino, N-benzyl-N-methylamino, 2-furylmethoxy,
- 5 3-furylmethoxy, 2-thienylmethoxy, 3-thienylmethoxy, 1-imidazolylmethoxy, 2-imidazolylmethoxy, 2-imidazol-1-ylethoxy, 3-imidazol-1-ylpropoxy, 2-imidazol-1-ylethylamino, 3-imidazol-1-ylpropylamino, 2-oxazolylmethoxy,
- 
- 4-oxazolylmethoxy, 5-oxazolylmethoxy, 2-thiazolylmethoxy, 4-thiazolylmethoxy, 5-thiazolylmethoxy, 1,2,3-triazol-1-ylmethoxy, 2-(1,2,3-triazol-1-yl)ethoxy,
- 10 3-(1,2,3-triazol-1-yl)propoxy, 1,2,3-triazol-1-ylmethylamino, 2-(1,2,3-triazol-1-yl)ethylamino, 3-(1,2,3-triazol-1-yl)propylamino, 1,2,4-triazol-1-ylmethoxy, 2-(1,2,4-triazol-1-yl)ethoxy, 3-(1,2,4-triazol-1-yl)propoxy, 1,2,4-triazol-1-ylmethylamino, 2-(1,2,4-triazol-1-yl)ethylamino, 3-(1,2,4-triazol-1-yl)propylamino, 2-pyridylmethoxy, 3-pyridylmethoxy, 4-pyridylmethoxy, 2-pyrid-2-ylethoxy, 2-pyrid-3-ylethoxy, 2-pyrid-4-ylethoxy, 3-pyrid-2-ylpropoxy,
- 15 3-pyrid-3-ylpropoxy, 3-pyrid-4-ylpropoxy, pyrrolidin-1-yl, pyrrolidin-3-yloxy, pyrrolidin-3-ylamino, N-methyl-N-(3-pyrrolidinyl)amino, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylamino, piperidin-4-ylamino, N-methyl-N-(3-piperidinyl)amino, N-methyl-N-(4-piperidinyl)amino, homopiperidin-1-yl, homopiperidin-3-yloxy,
- 20 homopiperidin-4-yloxy, piperazin-1-yl, homopiperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino, pyrrolidin-3-ylmethoxy, 2-pyrrolidin-3-ylethoxy, 3-pyrrolidin-3-ylpropoxy,
- 25 pyrrolidin-3-ylmethylamino, 2-pyrrolidin-3-ylethylamino, 3-pyrrolidin-3-ylpropylamino, 2-imidazolidin-1-ylethoxy, 3-imidazolidin-1-ylpropoxy, imidazolidin-2-ylmethoxy, 2-imidazolidin-2-ylethoxy, 3-imidazolidin-2-ylpropoxy, 2-imidazolidin-1-ylethylamino, 3-imidazolidin-1-ylpropylamino, 2-imidazolidin-2-ylethylamino, 3-imidazolidin-2-ylpropylamino, 2-morpholinoethoxy, 3-morpholinopropoxy,
- 30 2-morpholinoethylamino, 3-morpholinopropylamino, morpholin-2-ylmethoxy, 2-morpholin-2-ylethoxy, 3-morpholin-2-ylpropoxy, 2-morpholin-2-ylethylamino, 3-morpholin-2-ylpropylamino, morpholin-3-ylmethoxy, 2-morpholin-3-ylethoxy,

- 3-morpholin-3-ylpropoxy, 2-morpholin-3-ylethylamino, 3-morpholin-3-ylpropylamino, 2-(tetrahydro-4H-1,4-thiazin-4-yl)ethoxy, 3-(tetrahydro-4H-1,4-thiazin-4-yl)propoxy, 2-(tetrahydro-4H-1,4-thiazin-4-yl)ethylamino, 3-(tetrahydro-4H-1,4-thiazin-4-yl)propylamino, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethoxy, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propoxy, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethylamino, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propylamino, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidinoethylamino, 3-piperidinopropylamino, piperidin-2-ylmethoxy, 2-piperidin-2-ylethoxy, 3-piperidin-2-ylpropoxy, piperidin-2-ylmethylamino, 2-piperidin-2-ylethylamino, 3-piperidin-2-ylpropylamino, piperidin-3-ylmethoxy, 10 2-piperidin-3-ylethoxy, 3-piperidin-3-ylpropoxy, piperidin-3-ylmethylamino, 2-piperidin-3-ylethylamino, 3-piperidin-3-ylpropylamino, piperidin-4-ylmethoxy, 2-piperidin-4-ylethoxy, 3-piperidin-4-ylpropoxy, piperidin-4-ylmethylamino, 2-piperidin-4-ylethylamino, 3-piperidin-4-ylpropylamino, 2-homopiperidin-1-ylethoxy, 3-homopiperidin-1-ylpropoxy, 2-homopiperidin-1-ylethylamino, 15 3-homopiperidin-1-ylpropylamino, homopiperidin-2-ylmethoxy, homopiperidin-2-ylmethylamino, homopiperidin-3-ylmethoxy, homopiperidin-3-ylmethylamino, homopiperidin-4-ylmethoxy, homopiperidin-4-ylmethylamino, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino, piperazin-2-ylmethoxy, 20 piperazin-2-ylmethylamino, 2-piperazin-2-ylethoxy, 3-piperazin-2-ylpropoxy, 2-piperazin-2-ylethylamino, 3-piperazin-2-ylpropylamino, 2-homopiperazin-1-ylethoxy, 3-homopiperazin-1-ylpropoxy, 2-homopiperazin-1-ylethylamino, 3-homopiperazin-1-ylpropylamino, homopiperazin-2-ylmethoxy and homopiperazin-2-ylmethylamino, 25 and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a R<sup>1</sup> substituent is optionally separated by the insertion of an O or NH group, and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy, 30 ethoxy, propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl, and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2 oxo substituents,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and diethylamino;

5 (e) n is 0;

(f) R<sup>3</sup> is (1-6C)alkyl;

(g) R<sup>3</sup> is methyl;

(h) p is 0;

(i) Q<sup>1</sup> is aryl or heteroaryl and Q<sup>1</sup> is optionally substituted with 1, 2 or 3 substituents,  
 10 which may be the same or different, selected from hydroxy, halogeno, trifluoromethyl, cyano, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :

15 
$$-X^3-Q^4$$


wherein X<sup>3</sup> is a direct bond or is selected from O, NH and N(Me) and Q<sup>4</sup> is aryl, heteroaryl or heterocyclyl, and any Q<sup>4</sup> group optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, hydroxy, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

20 and wherein any Q<sup>4</sup> group when it is heterocyclyl optionally bears 1 or 2 oxo substituents,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a Q<sup>1</sup> group optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group one or more (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

25 (j) Q<sup>1</sup> is a phenyl group or a heteroaryl group comprising an aromatic 5- or 6-membered monocyclic ring, a 9- or 10-membered bicyclic ring or a 13- or 14-membered tricyclic ring each containing 1 oxygen heteroatom or 1 or 2 nitrogen heteroatoms and optionally containing a further heteroatom selected from nitrogen, oxygen and sulphur, and Q<sup>1</sup> is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from  
 30 hydroxy, halogeno, trifluoromethyl, cyano, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,

N-(1-6C)alkyl-(2-6C)alkanoylamino, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein  $X^3$  is a direct bond or is selected from O, NH and N(Me) and  $Q^4$  is phenyl, a

- 5 heteroaryl group comprising an aromatic 5- or 6-membered monocyclic ring containing 1 oxygen heteroatom or 1 or 2 nitrogen heteroatoms and optionally containing a further heteroatom selected from nitrogen, oxygen and sulphur or a heterocyclyl group comprising a non-aromatic saturated or partially saturated 3- to 10-membered monocyclic ring with up to 3 heteroatoms selected from oxygen, nitrogen and sulphur, and any  $Q^4$  group optionally bears
- 10 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, hydroxy, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any  $Q^4$  group when it is a heterocyclyl group optionally bears 1 or 2 oxo substituents,

- 15 and wherein any  $CH_2$  or  $CH_3$  group within a  $Q^1$  group optionally bears on each said  $CH_2$  or  $CH_3$  group one or more (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

- (k)  $Q^1$  is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano, amino,
- 20 methyl, ethyl, propyl, vinyl, ethynyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino, N-ethyl-N-methylamino, N-methyl-N-propylamino, acetoxy, acetamido, propionamido, acrylamido, propiolamido, N-methylacetamido, methanesulphonylamino, ethanesulphonylamino and N-methylmethanesulphonylamino, or from a group of the formula :

- 25  $-X^3-Q^4$

wherein  $X^3$  is a direct bond or is selected from O, NH and N(Me) and  $Q^4$  is phenyl, furyl, thienyl, imidazolyl, oxazolyl, thiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, azetidiny, tetrahydrofuranyl, pyrrolinyl, pyrrolidinyl, morpholinyl, tetrahydro-

- 30 piperazinyl or homopiperazinyl, and any  $Q^4$  group optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, trifluoromethyl, hydroxy,



amino, methyl, ethyl, propyl, vinyl, ethynyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino and diethylamino,


and wherein any  $Q^4$  group when it is a heterocyclyl group optionally bears 1 or 2 oxo substituents,

- 5 and wherein any  $CH_2$  or  $CH_3$  group within a  $Q^1$  group optionally bears on each said  $CH_2$  or  $CH_3$  group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino,

---

dimethylamino and diethylamino;

- (l)  $Q^1$  is optionally substituted with 1, 2 or 3 substituents, which may be the same or  
 10 different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano, amino, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino, N-ethyl-N-methylamino, N-methyl-N-propylamino, acetamido, N-methylacetamido, methanesulphonylamino, ethanesulphonylamino, N-methylmethanesulphonylamino, 1-azetidiny, 15 2- or 3-tetrahydrofuranyl, 3-pyrrolin-1-yl, 1-pyrrolidinyl, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, 1-piperidinyl, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or from a group of the formula :



wherein  $X^3$  is a direct bond or is selected from O, NH and N(Me) and  $Q^4$  is phenyl,

- 20 2- or 3-furyl, 2- or 3-thienyl, 1- or 2-imidazolyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 2-, 3- or 4-pyridyl, 3- or 4-pyridazinyl, 2- or 4-pyrimidinyl or 2-pyrazinyl, and any  $Q^4$  group optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, trifluoromethyl, hydroxy, amino, methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino,

- 25 and wherein any heterocyclyl group within  $Q^1$  optionally bears 1 or 2 oxo substituents.

and wherein any  $CH_2$  or  $CH_3$  group within a  $Q^1$  group optionally bears on each said  $CH_2$  or  $CH_3$  group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino;

- (m)  $Q^1$  is optionally substituted with 1, 2 or 3 substituents, which may be the same or  
 30 different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano, amino, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino, N-ethyl-N-methylamino,

N-methyl-N-propylamino, acetamido, N-methylacetamido, methanesulphonylamino, ethanesulphonylamino, N-methylmethanesulphonylamino, 1-azetidiny, 2- or 3-tetrahydrofuranyl, 3-pyrrolin-1-yl, 1-pyrrolidinyl, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, 1-piperidinyl,

5 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl,

and wherein any heterocyclyl group within Q<sup>1</sup> optionally bears 1 or 2 oxo substituents, and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a Q<sup>1</sup> group optionally bears on each said

---

CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino;

10 (n) Q<sup>1</sup> is a phenyl group which is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from those defined in paragraphs (k) to (m) hereinbefore;

(o) Q<sup>1</sup> is a 5- or 6-membered monocyclic heteroaryl ring containing 1 oxygen heteroatom or 1 or 2 nitrogen heteroatoms and optionally containing a further heteroatom selected from  
15 nitrogen, oxygen and sulphur, and Q<sup>1</sup> is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from those defined in paragraphs (k) to (m) hereinbefore;

(p) Q<sup>1</sup> is a 9- or 10-membered bicyclic heteroaryl ring containing 1 oxygen heteroatom or 1 or 2 nitrogen heteroatoms and optionally containing a further heteroatom selected from  
20 nitrogen, oxygen and sulphur, and Q<sup>1</sup> is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from those defined in paragraphs (k) to (m) hereinbefore;

(q) Q<sup>1</sup> is a 13- or 14-membered tricyclic heteroaryl ring containing 1 oxygen heteroatom or 1 or 2 nitrogen heteroatoms and optionally containing a further heteroatom selected from  
25 nitrogen, oxygen and sulphur, and Q<sup>1</sup> is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, halogeno, trifluoromethyl, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

(r) Q<sup>1</sup> is phenyl, furyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, benzofuranyl, indolyl,  
30 benzothiophenyl, benzoxazolyl, benzimidazolyl, benzothiazolyl, indazolyl, benzofurazanyl, quinolyl, isoquinolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, carbazolyl, dibenzofuranyl,

dibenzothiophenyl or xanthenyl which optionally bears 1 or 2 substituents selected from those defined in paragraphs (k) to (m) hereinbefore;

(s)  $Q^1$  is 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-imidazolyl, 3- or 4-pyrazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl,

5 2-, 3- or 4-pyridyl, 3- or 4-pyridazinyl, 2-, 4- or 5-pyrimidinyl or 2-pyrazinyl which optionally bears 1 or 2 substituents selected from those defined in paragraphs (l) and (m) hereinbefore;

(t)  $Q^1$  is 2-, 3-, 5- or 6-benzofuranyl, 2-, 3-, 5- or 6-indolyl,

2-, 3-, 5- or 6-benzothiophenyl, 2-, 5- or 6-benzoxazolyl, 2-, 5- or 6-benzimidazolyl,

2-, 5- or 6-benzothiazolyl, 3-, 5- or 6-indazolyl, 5-benzofurazanyl, 2-, 3-, 6- or 7-quinolyl,

10 3-, 6- or 7-isoquinolyl, 2-, 6- or 7-quinazolinyl, 2-, 6- or 7-quinoxaliny, or

1,8-naphthyridin-2-yl or 1,8-naphthyridin-3-yl which optionally bears 1 or 2 substituents selected from those defined in paragraphs (l) and (m) hereinbefore; and

(u)  $Q^1$  is 1-, 2- or 3-carbazolyl, 1-, 2-, 3- or 4-dibenzofuranyl or 1-, 2-, 3- or


4-dibenzothiophenyl which optionally bears 1 or 2 substituents selected from those defined in

15 paragraph (q) hereinbefore;

A preferred compound of the invention is a pyrimidine derivative of the Formula I wherein m is 0, 1, 2 or 3, and each  $R^1$  group, which may be the same or different, is selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylthio, methylsulphiny, methylsulphonyl,

20 methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, propargylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino,

N-methylcarbamoyl, N,N-dimethylcarbamoyl and acetamido, or from a group of the formula :



wherein  $X^1$  is a direct bond or is selected from O, NH and N(Me) and  $Q^2$  is benzyl,

25 2-furylmethyl, 3-furylmethyl, 2-thienylmethyl, 3-thienylmethyl,

1-imidazolylmethyl, 2-imidazolylmethyl, 2-imidazol-1-ylethyl, 3-imidazol-1-ylpropyl,

4-imidazol-1-ylbutyl, 2-oxazolylmethyl, 4-oxazolylmethyl, 5-oxazolylmethyl,

2-thiazolylmethyl, 4-thiazolylmethyl, 5-thiazolylmethyl, 1,2,3-triazol-1-ylmethyl,

2-(1,2,3-triazol-1-yl)ethyl, 3-(1,2,3-triazol-1-yl)propyl, 1,2,4-triazol-1-ylmethyl,

30 2-(1,2,4-triazol-1-yl)ethyl, 3-(1,2,4-triazol-1-yl)propyl, 2-pyridylmethyl, 3-pyridylmethyl,

4-pyridylmethyl, 2-pyrid-2-ylethyl, 2-pyrid-3-ylethyl, 2-pyrid-4-ylethyl, 3-pyrid-2-ylpropyl,

3-pyrid-3-ylpropyl, 3-pyrid-4-ylpropyl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl,

- morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, homopiperidin-1-yl, homopiperidin-2-yl, homopiperidin-3-yl, homopiperidin-4-yl, piperazin-1-yl, homopiperazin-1-yl, pyrrolidin-1-ylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-1-ylpropyl, 5 pyrrolidin-2-ylmethyl, 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-2-ylpropyl, pyrrolidin-3-ylmethyl, 2-pyrrolidin-3-ylethyl, 3-pyrrolidin-3-ylpropyl, imidazolidin-1-ylmethyl, 2-imidazolidin-1-ylethyl, 3-imidazolidin-1-ylpropyl, imidazolidin-2-ylmethyl, 2-imidazolidin-2-ylethyl, 3-imidazolidin-2-ylpropyl, morpholinomethyl, 2-morpholinoethyl, 3-morpholinopropyl, morpholin-2-ylmethyl, 10 2-morpholin-2-ylethyl, 3-morpholin-2-ylpropyl, morpholin-3-ylmethyl, 2-morpholin-3-ylethyl, 3-morpholin-3-ylpropyl, tetrahydro-4H-1,4-thiazin-4-ylmethyl, 2-(tetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(tetrahydro-4H-1,4-thiazin-4-yl)propyl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-ylmethyl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propyl, piperidinomethyl, 15 2-piperidinoethyl, 3-piperidinopropyl, piperidin-2-ylmethyl, 2-piperidin-2-ylethyl, 3-piperidin-2-ylpropyl, piperidin-3-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-3-ylpropyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, homopiperidin-1-ylmethyl, 2-homopiperidin-1-ylethyl, 3-homopiperidin-1-ylpropyl, homopiperidin-2-ylmethyl, 2-homopiperidin-2-ylethyl, 3-homopiperidin-2-ylpropyl, 20 homopiperidin-3-ylmethyl, 2-homopiperidin-3-ylethyl, 3-homopiperidin-3-ylpropyl, homopiperidin-4-ylmethyl, 2-homopiperidin-4-ylethyl, 3-homopiperidin-4-ylpropyl, piperazin-1-ylmethyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, piperazin-2-ylmethyl, 2-piperazin-2-ylethyl, 3-piperazin-2-ylpropyl, homopiperazin-1-ylmethyl, 2-homopiperazin-1-ylethyl, 3-homopiperazin-1-ylpropyl, homopiperazin-2-ylmethyl, 25 2-homopiperazin-2-ylethyl or 3-homopiperazin-2-ylpropyl,
- and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a R<sup>1</sup> substituent is optionally separated by the insertion of a group selected from O and NH,
- and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from 30 hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,


and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2 oxo substituents,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and diethylamino;

each of n and p is 0;

R<sup>3</sup> is hydrogen or methyl; and

Q<sup>1</sup> is phenyl, 2-furyl, 2-thienyl, 4-oxazolyl, 5-isoxazolyl, 2- or 4-imidazolyl, 3- or 4-pyrazolyl, 4-thiazolyl, 5-isothiazolyl, 2-, 3- or 4-pyridyl, 4-pyridazinyl, 4- or 5-pyrimidinyl, 2- or 6-benzofuranyl, 2- or 6-indolyl, 2- or 6-benzothiophenyl, 2- or 6-quinolyl or 2- or 4-dibenzofuranyl which optionally bears 1, 2 or 3 substituents, , which may be the same or different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano, amino, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino, N-ethyl-N-methylamino, N-methyl-N-propylamino, acetamido, N-methylacetamido, methanesulphonylamino, ethanesulphonylamino, N-methylmethanesulphonylamino, 1-azetidiny, 3-pyrrolin-1-yl, 1-pyrrolidinyl, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, 1-piperidinyl, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or from a group of the formula :



wherein X<sup>3</sup> is a direct bond or is selected from O, NH and N(Me) and Q<sup>4</sup> is phenyl, 2- or 3-furyl, 2- or 3-thienyl, 1- or 2-imidazolyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 2-, 3- or 4-pyridyl, 3- or 4-pyridazinyl, 2- or 4-pyrimidinyl or 2-pyrazinyl, and any Q<sup>4</sup> group optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, trifluoromethyl, hydroxy, amino, methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino,

and wherein any heterocyclyl group within Q<sup>1</sup> optionally bears 1 or 2 oxo substituents,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a Q<sup>1</sup> group optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino; or a pharmaceutically acceptable salt or in-vivo-cleavable ester thereof.

A further preferred compound of the invention is a pyrimidine derivative of the Formula I

- wherein m is 0, 1 or 2 and each R<sup>1</sup> group, which may be the same or different, is selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylthio, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl, acetamido, benzyloxy, benzylamino, N-benzyl-N-methylamino, 2-furylmethoxy, 3-furylmethoxy, 2-imidazol-1-ylethylamino, 3-imidazol-1-ylpropylamino, 2-(1,2,4-triazol-1-yl)ethylamino, 3-(1,2,4-triazol-1-yl)propylamino, 2-pyridylmethoxy, 3-pyridylmethoxy, 4-pyridylmethoxy, pyrrolidin-1-yl, pyrrolidin-3-yloxy, pyrrolidin-3-ylamino, N-methyl-N-(3-pyrrolidinyl)amino, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylamino, piperidin-4-ylamino, N-methyl-N-(3-piperidinyl)amino, N-methyl-N-(4-piperidinyl)amino, homopiperidin-1-yl, homopiperidin-3-yloxy, homopiperidin-4-yloxy, piperazin-1-yl, homopiperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino, pyrrolidin-3-ylmethoxy, 2-pyrrolidin-3-ylethoxy, 3-pyrrolidin-3-ylpropoxy, pyrrolidin-3-ylmethylamino, 2-pyrrolidin-3-ylethylamino, 3-pyrrolidin-3-ylpropylamino, 2-imidazolidin-1-ylethoxy, 3-imidazolidin-1-ylpropoxy, imidazolidin-2-ylmethoxy, 2-imidazolidin-2-ylethoxy, 3-imidazolidin-2-ylpropoxy, 2-imidazolidin-1-ylethylamino, 3-imidazolidin-1-ylpropylamino, 2-imidazolidin-2-ylethylamino, 3-imidazolidin-2-ylpropylamino, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-morpholinoethylamino, 3-morpholinopropylamino, morpholin-2-ylmethoxy, 2-morpholin-2-ylethoxy, 3-morpholin-2-ylpropoxy, 2-morpholin-2-ylethylamino, 3-morpholin-2-ylpropylamino, morpholin-3-ylmethoxy, 2-morpholin-3-ylethoxy, 3-morpholin-3-ylpropoxy, 2-morpholin-3-ylethylamino, 3-morpholin-3-ylpropylamino, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidinoethylamino, 3-piperidinopropylamino, piperidin-2-ylmethoxy, 2-piperidin-2-ylethoxy, 3-piperidin-2-ylpropoxy, piperidin-2-ylmethylamino, 2-piperidin-2-ylethylamino, 3-piperidin-2-ylpropylamino, piperidin-3-ylmethoxy, 2-piperidin-3-ylethoxy, 3-piperidin-3-ylpropoxy,

piperidin-3-ylmethylamino, 2-piperidin-3-ylethylamino, 3-piperidin-3-ylpropylamino,  
 piperidin-4-ylmethoxy, 2-piperidin-4-ylethoxy, 3-piperidin-4-ylpropoxy,  
 piperidin-4-ylmethylamino, 2-piperidin-4-ylethylamino, 3-piperidin-4-ylpropylamino,  
 2-homopiperidin-1-ylethoxy, 3-homopiperidin-1-ylpropoxy, 2-homopiperidin-1-ylethylamino,  
 5 3-homopiperidin-1-ylpropylamino, homopiperidin-2-ylmethoxy,  
 homopiperidin-2-ylmethylamino, homopiperidin-3-ylmethoxy,  
 homopiperidin-3-ylmethylamino, homopiperidin-4-ylmethoxy,

---

homopiperidin-4-ylmethylamino, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy,  
 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino, piperazin-2-ylmethoxy,  
 10 piperazin-2-ylmethylamino, 2-piperazin-2-ylethoxy, 3-piperazin-2-ylpropoxy,  
 2-piperazin-2-ylethylamino, 3-piperazin-2-ylpropylamino, 2-homopiperazin-1-ylethoxy,  
 3-homopiperazin-1-ylpropoxy, 2-homopiperazin-1-ylethylamino,  
 3-homopiperazin-1-ylpropylamino, homopiperazin-2-ylmethoxy or  
 homopiperazin-2-ylmethylamino,

15 and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R<sup>1</sup>  
 optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from  
 hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy,  
 ethoxy propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,  
 and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2  
 20 oxo substituents,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each  
 said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy,  
 amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and  
 diethylamino;

25 each of n and p is 0;

R<sup>3</sup> is methyl; and

Q<sup>1</sup> is phenyl or 3- or 4-pyridyl which optionally bears 1, 2 or 3 substituents, which may be the  
 same or different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano,  
 amino, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylamino, ethylamino,  
 30 propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino,  
N-ethyl-N-methylamino, N-methyl-N-propylamino, acetamido, N-methylacetamido,  
 methanesulphonylamino, ethanesulphonylamino, N-methylmethanesulphonylamino,

1-azetidiny, 3-pyrrolin-1-yl, 1-pyrrolidinyl, morpholino, 1-piperidinyl, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a Q<sup>1</sup> group optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy; methylamino, ethylamino, dimethylamino and diethylamino;  
 5 or a pharmaceutically acceptable salt or in-vivo-cleavable ester thereof.

A more preferred compound of the invention is a pyrimidine derivative of the

---

Formula I

wherein m is 0, 1 or 2 and each R<sup>1</sup> group, which may be the same or different, is selected from  
 10 fluoro, chloro, bromo, amino, carbamoyl, methoxy, ethoxy, propoxy, methylthio, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, pyrrolidin-3-yloxy, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylamino, piperidin-4-ylamino, N-methyl-N-(3-piperidinyl)amino,  
 15 N-methyl-N-(4-piperidinyl)amino, homopiperidin-1-yl, homopiperidin-3-yloxy, homopiperidin-4-yloxy, piperazin-1-yl, homopiperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino,  
 20 pyrrolidin-3-ylmethoxy, 2-pyrrolidin-3-ylethoxy, 3-pyrrolidin-3-ylpropoxy, pyrrolidin-3-ylmethylamino, 2-pyrrolidin-3-ylethylamino, 3-pyrrolidin-3-ylpropylamino, 2-imidazolidin-1-ylethylamino, 3-imidazolidin-1-ylpropylamino, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-morpholinoethylamino, 3-morpholinopropylamino, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidinoethylamino, 3-piperidinopropylamino,  
 25 piperidin-3-ylmethoxy, piperidin-4-ylmethoxy, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino or piperazin-2-ylmethoxy,

and wherein any heterocyclyl group within a substituent on R<sup>1</sup> optionally bears 1 or 2 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro,  
 30 methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,



and wherein any heterocyclcyl group within a substituent on R<sup>1</sup> optionally bears an oxo substituent,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a R<sup>1</sup> substituent optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methylamino, ethylamino, dimethylamino and diethylamino;  
 5 each of n and p is 0;  
 R<sup>3</sup> is methyl; and

---

Q<sup>1</sup> is phenyl or 4-pyridyl which optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, amino, methyl, methoxy,  
 10 methylamino, ethylamino, dimethylamino, diethylamino, 1-pyrrolidinyl, morpholino, piperidino, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl,

and wherein any CH<sub>2</sub> or CH<sub>3</sub> group within a Q<sup>1</sup> group optionally bears on each said CH<sub>2</sub> or CH<sub>3</sub> group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino;  
 15 or a pharmaceutically acceptable salt or in-vivo-cleavable ester thereof.

A further more preferred compound of the invention is a pyrimidine derivative of the Formula I

wherein m is 1 or 2 and each R<sup>1</sup> group, which may be the same or different, is selected from chloro, carbamoyl, 3-dimethylaminopropoxy, 3-dimethylamino-2,2-dimethylpropoxy,  
 20 methylthio, 3-diethylaminopropylamino, 3-dimethylamino-2,2-dimethylpropylamino, 3-dimethylamino-2-hydroxypropylamino, N-isopropylpyrrolidin-3-yloxy, piperidin-4-yloxy, N-methylpiperidin-4-yloxy, N-ethylpiperidin-3-ylamino, N-methylpiperidin-4-ylamino, N-methyl-N-(N-methylpiperidin-4-yl)amino, 3-pyrrolidin-1-ylpropylamino, N-methylpyrrolidin-2-ylmethoxy, 2-(N-methylpyrrolidin-2-yl)ethoxy,  
 25 2-(N-methylpyrrolidin-2-yl)ethylamino, N-methylpiperidin-3-ylmethoxy and N,N'-dimethylpiperazin-2-ylmethoxy;

each of n and p is 0;

R<sup>3</sup> is methyl; and

Q<sup>1</sup> is phenyl which bears 1 or 2 substituents, which may be the same or different, selected  
 30 from fluoro, trifluoromethyl, dimethylamino, 1-pyrrolidinyl, morpholino, piperidino, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or Q<sup>1</sup> is 4-pyridyl which bears 1

substituent selected from dimethylamino, 1-pyrrolidinyl, morpholino, piperidino, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or a pharmaceutically acceptable salt or in-vivo-cleavable ester thereof.

A particular preferred compound of the invention is, for example :-


- 5 6-carbamoyl-2-chloro-4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]pyrimidine,  
4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]-2-methylthiopyrimidine,  
2-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine,  
4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-(N-methylpiperidin-  
4-yloxy)pyrimidine,
- 10 2-(3-dimethylaminopropoxy)-4-[2-methyl-5-(2-morpholinopyrid-  
4-ylcarbonylamino)anilino]pyrimidine,  
2-(3-dimethylamino-2,2-dimethylpropoxy)-4-[2-methyl-5-(2-morpholinopyrid-  
4-ylcarbonylamino)anilino]pyrimidine,  
4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-(N-methylpiperidin-
- 15 3-ylmethoxy)pyrimidine,  
2-[N-methyl-N-(N-methylpiperidin-4-yl)amino]-4-[2-methyl-5-(2-morpholinopyrid-  
4-ylcarbonylamino)anilino]pyrimidine and  
4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-[2-(N-methylpyrrolidin-  
2-yl)ethylamino]pyrimidine;
- 20 or a pharmaceutically acceptable salt or in-vivo-cleavable ester thereof.

A pyrimidine derivative of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes, when used to prepare a novel pyrimidine derivative of the Formula I are provided as a further feature of the invention and

25 are illustrated by the following representative process variants in which, unless otherwise stated,  $R^1$ ,  $m$ ,  $R^2$ ,  $n$ ,  $R^3$ ,  $p$  and  $Q^1$  have any of the meanings defined hereinbefore. Necessary starting materials may be obtained by standard procedures of organic chemistry. The preparation of such starting materials is described in conjunction with the following representative process variants and within the accompanying Examples. Alternatively


30 necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

(a) A compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, may be prepared by reacting an aniline of the Formula II



II

with an acid of the Formula III, or a reactive derivative thereof,



5

under standard amide bond forming conditions, wherein variable groups are as defined hereinbefore and wherein any functional groups are protected if necessary, and

- (i) removing any protecting groups; and
- (ii) optionally forming a pharmaceutically-acceptable salt or in-vivo-cleavable ester.

10

A suitable activated derivative of an acid of the Formula III is, for example, an acyl halide, for example an acyl chloride formed by the reaction of the acid and an inorganic acid chloride, for example thionyl chloride; a mixed anhydride, for example an anhydride formed by the reaction of the acid and a chloroformate such as isobutyl chloroformate; an active ester, for example an ester formed by the reaction of the acid with a phenol such as pentafluorophenol, with an ester such as pentafluorophenyl trifluoroacetate or with an alcohol such as N-hydroxybenzotriazole; an acyl azide, for example an azide formed by the reaction of the acid and an azide such as diphenylphosphoryl azide; an acyl cyanide, for example a cyanide formed by the reaction of an acid and a cyanide such as diethylphosphoryl cyanide; or the product of the reaction of the acid and a carbodiimide such as dicyclohexylcarbodiimide.

20

The reaction is preferably carried out in the presence of a suitable base such as, for example, an alkali or alkaline earth metal carbonate, alkoxide, hydroxide or hydride, for example sodium carbonate, potassium carbonate, sodium ethoxide, potassium butoxide, sodium hydroxide, potassium hydroxide, sodium hydride or potassium hydride, or an organometallic base such as an alkyl-lithium, for example n-butyl-lithium, or a dialkylamino-lithium, for example lithium di-isopropylamide, or, for example, an organic

25

amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, morpholine or diazabicyclo[5.4.0]undec-7-ene. The reaction is also preferably carried out in a suitable inert solvent or diluent, for example tetrahydrofuran, methylene chloride, 1,2-dimethoxyethane, N,N-dimethylformamide, N,N-dimethylacetamide, 5 N-methylpyrrolidin-2-one, dimethylsulphoxide or acetone, and at a temperature in the range, for example, -78 to 150°C, conveniently at or near ambient temperature.

Typically a carbodiimide coupling reagent is used in the presence of an organic solvent (preferably an anhydrous polar aprotic organic solvent) at a non-extreme temperature, for example in the region -10 to 40°C, typically at ambient temperature of about 20°C.

10 Protecting groups may in general be chosen from any of the groups described in the literature or known to the skilled chemist as appropriate for the protection of the group in question and may be introduced by conventional methods. Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so 15 as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Specific examples of protecting groups are given below for the sake of convenience, in which "lower", as in, for example, lower alkyl, signifies that the group to which it is applied preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive.

20 Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned is of course within the scope of the invention.

A carboxy protecting group may be the residue of an ester-forming aliphatic or arylaliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably 25 containing 1-20 carbon atoms). Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (for example isopropyl, tert-butyl); lower alkoxy lower alkyl groups (for example methoxymethyl, ethoxymethyl, isobutoxymethyl); lower aliphatic acyloxy lower alkyl groups, (for example acetoxymethyl, propionyloxymethyl, butyryloxymethyl, pivaloyloxymethyl); lower alkoxycarbonyloxy lower alkyl groups (for 30 example 1-methoxycarbonyloxyethyl, 1-ethoxycarbonyloxyethyl); aryl lower alkyl groups (for example benzyl, p-methoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (for example trimethylsilyl and tert-butyldimethylsilyl); tri(lower

alkyl)silyl lower alkyl groups (for example trimethylsilylethyl); and (2-6C)alkenyl groups (for example allyl and vinyllethyl). Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, base-, metal- or enzymically-catalysed hydrolysis.


- 5        Examples of hydroxy protecting groups include lower alkyl groups (for example tert-butyl), lower alkenyl groups (for example allyl); lower alkanoyl groups (for example acetyl); lower alkoxycarbonyl groups (for example tert-butoxycarbonyl); lower
- 
- alkenyloxycarbonyl groups (for example allyloxycarbonyl); aryl lower alkoxycarbonyl groups (for example benzoyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, *o*-nitrobenzyloxycarbonyl,
- 10 *p*-nitrobenzyloxycarbonyl); tri lower alkylsilyl (for example trimethylsilyl, tert-butyldimethylsilyl) and aryl lower alkyl (for example benzyl) groups.

- Examples of amino protecting groups include formyl, aralkyl groups (for example benzyl and substituted benzyl, *p*-methoxybenzyl, nitrobenzyl and 2,4-dimethoxybenzyl, and triphenylmethyl); di-*p*-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (for
- 15 example tert-butoxycarbonyl); lower alkenyloxycarbonyl (for example allyloxycarbonyl); aryl lower alkoxycarbonyl groups (for example benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl; trialkylsilyl (for example trimethylsilyl and tert-butyldimethylsilyl); alkylidene (for example methylenidene); benzylenidene and substituted benzylenidene groups.

- 20        Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base-, metal- or enzymically-catalysed hydrolysis for groups such as *p*-nitrobenzyloxycarbonyl, hydrogenation for groups such as benzyl and photolytically for groups such as *o*-nitrobenzyloxycarbonyl.

- The reader is referred to Advanced Organic Chemistry, 4th Edition, by Jerry March,
- 25 published by John Wiley & Sons 1992, for general guidance on reaction conditions and reagents. The reader is referred to Protective Groups in Organic Synthesis, 2nd Edition, by Green *et al.*, published by John Wiley & Sons for general guidance on protecting groups.


The aniline of Formula II may be prepared by reduction of the corresponding nitro compound of Formula IV.



IV


Typical reaction conditions include the use of ammonium formate in the presence of a catalyst (for example palladium-on-carbon) in the presence of an organic solvent (preferably a polar protic solvent), preferably with heating, for example to about 60°C. Any functional groups are protected and deprotected as necessary.

The nitro compound of Formula IV may be prepared by the reaction, conveniently in the presence of a suitable base as defined hereinbefore, of an activated heteroaryl compound of the Formula V



V

10 wherein L is a displaceable group as defined hereinafter with an aniline of the Formula VI




VI

A suitable displaceable group L is, for example, a halogeno group such as fluoro, chloro or bromo, an activated phenoxy group such as pentafluorophenoxy, a (1-6C)alkanesulphonyloxy group such as methanesulphonyloxy or an arylsulphonyloxy group such as 4-toluenesulphonyloxy.


The reaction is conveniently carried out in the presence of a suitable inert diluent or carrier as defined hereinbefore and at a temperature in the range, for example, 20 to 200°C, conveniently in the range 50 to 150°C.

(b) A compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, may be prepared by the reaction, conveniently in the presence of a suitable base as defined hereinbefore, of an activated heteroaryl compound of the Formula V



V

wherein L is a displaceable group as defined hereinbefore, with an aniline of the Formula VII




VII

wherein variable groups are as defined hereinbefore and wherein any functional group is  
5 protected if necessary, and

- (i) removing any protecting groups; and
- (ii) optionally forming a pharmaceutically-acceptable salt or in-vivo-cleavable ester.

The reaction is conveniently carried out in the presence of a suitable inert diluent or  
10 carrier as defined hereinbefore and at a temperature in the range, for example, 20 to 200°C, conveniently in the range 50 to 150°C.


The aniline of Formula VII may be prepared by reduction of the corresponding nitro compound of Formula VIII.



VIII


15 under typical reaction conditions such as catalytic reduction with hydrogen gas in the presence of a metallic catalyst, for example palladium-on-carbon, or the use of ammonium formate in the presence of a metallic catalyst, for example palladium-on-carbon.

The nitro compound of the Formula VIII may be prepared by the coupling of an aniline of the Formula IX



IX

with an acid of the Formula III, or a reactive derivative thereof as defined hereinbefore,



under standard amide bond forming conditions as defined hereinbefore.

- (c) A compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein  $\text{R}^1$  is an amino, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, substituted (1-6C)alkylamino, substituted di-[(1-6C)alkyl]amino or a N-linked heterocyclyl substituent may be prepared by the reaction, conveniently in the presence of a suitable base as defined hereinbefore, of an appropriate amine with a pyrimidine derivative of the Formula I wherein  $\text{R}^1$  is a suitable displaceable group as defined hereinbefore and wherein other variable groups are as defined hereinbefore and wherein any functional group is protected if necessary, and

- (i) removing any protecting groups; and
- (ii) optionally forming a pharmaceutically-acceptable salt or in-vivo-cleavable ester.

- 15 The reaction is conveniently carried out in the presence of a suitable inert diluent or carrier as defined hereinbefore and at a temperature in the range, for example, 20 to 200°C, conveniently in the range 75 to 150°C.

- (d) A compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein  $\text{R}^1$  is a (1-6C)alkoxy or substituted (1-6C)alkoxy substituent may be prepared by the reaction, conveniently in the presence of a suitable base as defined hereinbefore, of an appropriate alcohol with a pyrimidine derivative of the Formula I wherein  $\text{R}^1$  is a suitable displaceable group as defined hereinbefore and wherein other variable groups are as defined hereinbefore and wherein any functional group is protected if necessary, and

- (i) removing any protecting groups; and
- 25 (ii) optionally forming a pharmaceutically-acceptable salt or in-vivo-cleavable ester.

The reaction is conveniently carried out in the presence of a suitable inert diluent or carrier as defined hereinbefore and at a temperature in the range, for example, 20 to 200°C, conveniently in the range 100 to 180°C.

- 30 The following biological assays and Examples serve to illustrate the present invention.



**Biological Assays**

The following assays can be used to measure the p38 kinase-inhibitory, the TNF-inhibitory and anti-arthritis effects of the compounds of the present invention:

**In vitro enzyme assay**

- 5           The ability of compounds of the invention to inhibit the enzyme p38 kinase was assessed. Activity of test compounds against each of the p38 $\alpha$  and p38 $\beta$  isoforms of the enzyme was determined.

---

Human recombinant MKK6 (GenBank Accession Number G1209672) was isolated from Image clone 45578 (Genomics, 1996, 33, 151) and utilised to produce protein in the  
 10 form of a GST fusion protein in a pGEX vector using analogous procedures to those disclosed by J. Han *et al.*, Journal of Biological Chemistry, 1996, 271, 2886-2891. p38 $\alpha$  (GenBank Accession Number G529039) and p38 $\beta$  (GenBank Accession Number G1469305) were isolated by PCR amplification of human lymphoblastoid cDNA (GenBank Accession Number GM1416) and human foetal brain cDNA [synthesised from mRNA (Clontech, catalogue  
 15 no. 6525-1) using a Gibco superscript cDNA synthesis kit] respectively using oligonucleotides designed for the 5' and 3' ends of the human p38 $\alpha$  and p38 $\beta$  genes using analogous procedures to those described by J.Han *et al.*, Biochimica et Biophysica Acta, 1995, 1265, 224-227 and Y. Jiang *et al.*, Journal of Biological Chemistry, 1996, 271, 17920-17926.

- Both p38 protein isoforms were expressed in *e coli* in PET vectors. Human  
 20 recombinant p38 $\alpha$  and p38 $\beta$  isoforms were produced as 5' c-myc, 6His tagged proteins. Both MKK6 and the p38 proteins were purified using standard protocols: the GST MKK6 was purified using a glutathione sepharose column and the p38 proteins were purified using nickel chelate columns.

- The p38 enzymes were activated prior to use by incubation with MKK6 for 3 hours at  
 25 30°C. The unactivated coli-expressed MKK6 retained sufficient activity to fully activate both isoforms of p38. The activation incubate comprised p38 $\alpha$  (10 $\mu$ l of 10mg/ml) or p38 $\beta$  (10 $\mu$ l of 5mg/ml) together with MKK6 (10 $\mu$ l of 1mg/ml), 'Kinase buffer' [100 $\mu$ l; pH 7.4 buffer comprising Tris (50mM), EGTA (0.1mM), sodium orthovanadate (0.1mM) and  $\beta$ -mercaptoethanol (0.1%)] and MgATP (30 $\mu$ l of 50mM Mg(OCOCH<sub>3</sub>)<sub>2</sub> and 0.5mM ATP).  
 30 This produced enough activated p38 enzyme for 3 Microtiter plates.

Test compounds were solubilised in DMSO and 10µl of a 1:10 diluted sample in 'Kinase Buffer' was added to a well in a Microtiter plate. For single dose testing, the compounds were tested at 10µM. 'Kinase Assay Mix' [30µl; comprising Myelin Basic Protein (Gibco BRL cat. no. 1322B-010; 1ml of a 3.33mg/ml solution in water), activated p38 enzyme (50µl) and 'Kinase Buffer' (2ml)] was then added followed by 'Labelled ATP' [10µl; comprising 50µM ATP, 0.1µCi <sup>33</sup>P ATP (Amersham International cat. no. BF1000) and 50mM Mg(OCOCH<sub>3</sub>)<sub>2</sub>]. The plates were incubated at room temperature with gentle agitation.

Plates containing p38α were incubated for 90min and plates containing p38β were incubated for 45min. Incubation was stopped by the addition of 50µl of 20% trichloroacetic acid (TCA).  
10 The precipitated protein was phosphorylated by p38 kinase and test compounds were assessed for their ability to inhibit this phosphorylation. The plates were filtered using a Canberra Packard Unifilter and washed with 2% TCA, dried overnight and counted on a Top Count scintillation counter.

Test compounds were tested initially at a single dose and active compounds were  
15 retested to allow IC<sub>50</sub> values to be determined.

#### In vitro cell-based assays

##### (i) PBMC

The ability of compounds of this invention to inhibit TNFα production was assessed by using human peripheral blood mononuclear cells which synthesise and secrete TNFα when  
20 stimulated with lipopolysaccharide.

Peripheral blood mononuclear cells (PBMC) were isolated from heparinised (10units/ml heparin) human blood by density centrifugation (Lymphoprep<sup>TM</sup>; Nycomed). Mononuclear cells were resuspended in culture medium [RPMI 1640 medium (Gibco) supplemented with 50 units/ml penicillin, 50µg/ml streptomycin, 2mM glutamine and  
25 1% heat-inactivated human AB serum (Sigma H-1513)]. Compounds were solubilised in DMSO at a concentration of 50mM, diluted 1:100 in culture medium and subsequently serial dilutions were made in culture medium containing 1% DMSO. PBMCs (2.4x10<sup>5</sup> cells in 160µl culture medium) were incubated with 20µl of varying concentrations of test compound (triplicate cultures) or 20µl culture medium containing 1% DMSO (control wells) for  
30 30 minutes at 37°C in a humidified (5%CO<sub>2</sub>/95% air) incubator (Falcon 3072 ; 96 well flat-bottom tissue culture plates). 20µl lipopolysaccharide [LPS E.Coli 0111:B4 (Sigma

L-4130), final concentration 10µg/ml] solubilised in culture medium was added to appropriate wells. 20µl culture medium was added to "medium alone" control wells. Six "LPS alone" and four "medium alone" controls were included on each 96 well plate. Varying concentrations of a known TNFα inhibitor were included in each test, i.e. an inhibitor of the PDE Type IV enzyme (for example see Semmler, J. Wachtel. H and Endres, S., Int. J. Immunopharmac. (1993), 15(3), 409-413) or an inhibitor of proTNFα convertase (for example, see McGeehan, G. M. *et al.* Nature (1994) 370, 558-561). Plates were incubated for 7 hours at 37°C

(humidified incubator) after which 100µl of the supernatant was removed from each well and stored at -70°C (96 well round-bottom plates; Corning 25850). TNFα levels were determined in each sample using a human TNFα ELISA (see WO92/10190 and Current Protocols in Molecular Biology, vol 2 by Frederick M. Ausbel et al., John Wiley and Sons Inc.).

$$\% \text{ inhibition} = \frac{(\text{LPS alone} - \text{medium alone}) - (\text{test concentration} - \text{medium alone})}{(\text{LPS alone} - \text{medium alone})} \times 100$$

(ii) **Human Whole Blood**

The ability of the compounds of this invention to inhibit TNFα production was also assessed in a human whole blood assay. Human whole blood secretes TNFα when stimulated with LPS. This property of blood forms the basis of an assay which is used as a secondary test for compounds which profile as active in the PBMC test.

Heparinised (10 units/ml) human blood was obtained from volunteers. 160µl whole blood were added to 96 well round-bottom plates (Corning 25850). Compounds were solubilised and serially diluted in RPMI 1640 medium (Gibco) supplemented with 50 units/ml penicillin, 50µg/ml streptomycin and 2mM glutamine, as detailed above. 20µl of each test concentration was added to appropriate wells (triplicate cultures). 20µl of RPMI 1640 medium supplemented with antibiotics and glutamine was added to control wells. Plates were incubated for 30 minutes at 37°C (humidified incubator), prior to addition of 20µl LPS (final concentration 10µg/ml). RPMI 1640 medium was added to control wells. Six "LPS alone" and four "medium alone" controls were included on each plate. A known TNFα synthesis/secretion inhibitor was included in each test. Plates were incubated for 6 hours at 37°C (humidified incubator). Plates were centrifuged (2000rpm for 10 minutes) and 100µl plasma removed and stored at -70°C (Corning 25850 plates). TNFα levels were measured by ELISA (see WO92/10190 and Current Protocols in Molecular Biology, vol 2 by Frederick M.

Ausbel *et al.*, John Wiley and Sons Inc.). The paired antibodies that were used in the ELIZA were obtained from R&D Systems (catalogue nos. MAB610 anti-human TNF $\alpha$  coating antibody, BAF210 biotinylated anti-human TNF $\alpha$  detect antibody).

#### **Ex vivo / In vivo assessment**

- 5           The ability of the compounds of this invention as *ex vivo* TNF $\alpha$  inhibitors were assessed in the rat or mouse. Briefly, groups of male Wistar Alderley Park (AP) rats (180-210g) were dosed with compound (6 rats) or drug vehicle (10 rats) by the appropriate route, for example peroral (p.o.), intraperitoneal (i.p.) or subcutaneous (s.c.). Ninety minutes later rats were sacrificed using a rising concentration of CO<sub>2</sub> and bled out via the posterior vena
- 
- 10   cavae into 5 Units of sodium heparin/ml blood. Blood samples were immediately placed on ice and centrifuged at 2000 rpm for 10 min at 4°C and the harvested plasmas frozen at -20°C for subsequent assay of their effect on TNF $\alpha$  production by LPS-stimulated human blood. The rat plasma samples were thawed and 175 $\mu$ l of each sample was added to a set format pattern in a 96 well round bottom plate (Corning 25850). 50 $\mu$ l of heparinized human blood
- 15   was then added to each well, mixed and the plate was incubated for 30 min at 37°C (humidified incubator). LPS (25 $\mu$ l; final concentration 10 $\mu$ g/ml) was added to the wells and incubation continued for a further 5.5 hours. Control wells were incubated with 25 $\mu$ l of medium alone. Plates were then centrifuged for 10 min at 2000 rpm and 200 $\mu$ l of the supernatants were transferred to a 96 well plate and frozen at -20°C for subsequent analysis of
- 20   TNF concentration by ELISA.

Data analysis by dedicated software calculates for each compound/dose:

$$\% \text{ inhibition of TNF}\alpha = \frac{\text{Mean TNF}\alpha (\text{Controls}) - \text{Mean TNF}\alpha (\text{Treated})}{\text{Mean TNF}\alpha (\text{Controls})} \times 100$$

Alternatively, mice could be used instead of rats in the above procedure.

#### 25   **Test as anti-arthritic agent**

Activity of a compound as an anti-arthritic agent was tested as follows. Acid soluble native type II collagen was shown by Trentham *et al.* [1] to be arthritogenic in rats; it caused polyarthritis when administered in Freund's incomplete adjuvant. This is now known as collagen-induced arthritis (CIA) and similar conditions can be induced in mice and primates.

- 30   Recent studies have shown that anti-TNF monoclonal antibodies [2] and TNF receptor-IgG

fusion proteins [3] ameliorate established CIA indicating that TNF plays a key role in the pathophysiology of CIA. Moreover, the remarkable efficacy reported for anti-TNF monoclonal antibodies in recent rheumatoid arthritis clinical trials indicates that TNF plays a major role in this chronic inflammatory disease. Thus CIA in DBA/1 mice as described in  
 5 references 2 and 3 is a tertiary model which can be used to demonstrate the anti-arthritic activity of a compound. Also see reference 4.

1. Trentham, D.E. *et al.*, (1977) J. Exp. Med., 146, 857.

2. Williams, R.O. *et al.*, (1992) Proc. Natl. Acad. Sci., 89, 9784.

3. Williams, R.O. *et al.*, (1995) Immunology, 84, 433.

10 4 Badger, M. B. *et al.*, (1996) The Journal of Pharmacology and Experimental Therapeutics, 279, 1453-1461.

Although the pharmacological properties of the compounds of the Formula I vary with structural change as expected, in general a compound of the Formula I gives over 30% inhibition of p38 $\alpha$  and/or p38 $\beta$  at concentrations up to 10 $\mu$ M and over 30% inhibition in the  
 15 PBMC test at concentrations up to 50 $\mu$ M. No physiologically unacceptable toxicity was observed at the effective dose for compounds tested of the present invention.

By way of example :-

6-carbamoyl-2-chloro-4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]pyrimidine has an IC<sub>50</sub> of approximately 0.03 $\mu$ M against p38 $\alpha$  and an IC<sub>50</sub> of approximately 16 $\mu$ M in the  
 20 Human Whole Blood test.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a pyrimidine derivative of the Formula I, or a pharmaceutically-acceptable or in-vivo-cleavable ester thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

25 The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for  
 30 example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

- 5           The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 0.5 g of active agent compounded with an appropriate and convenient amount of excipients  
10 which may vary from about 5 to about 98 percent by weight of the total composition.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula I will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.

- 15           In using a compound of the Formula I for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body  
20 weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred, particularly in tablet form. Typically, unit dosage forms will contain about 1 mg to 500 mg of a compound of this invention.

- According to a further aspect of the invention there is provided a pyrimidine derivative  
25 of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

- According to a further aspect of the invention there is provided the use of a pyrimidine derivative of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment  
30 or medical conditions mediated by cytokines.

In a further aspect the present invention provides a method of treating diseases or medical conditions mediated by cytokines which comprises administering to a warm-blooded

animal an effective amount of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof.

In a further aspect the present invention provides the use of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, in the  
5 manufacture of a medicament for use in the treatment of diseases or medical conditions mediated by TNF, IL-1, IL-6 or IL-8.

In a further aspect the present invention provides a method of treating diseases or  
medical conditions mediated by TNF, IL-1, IL-6 or IL-8 which comprises administering to a  
warm-blooded animal an effective amount of a compound of the Formula I or a  
10 pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof.

In a further aspect the present invention provides the use of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof in the  
manufacture of a medicament for use in the treatment of diseases or medical conditions  
mediated by TNF.

15 In a further aspect the present invention provides a method of treating diseases or medical conditions mediated by TNF which comprises administering to a warm-blooded animal an effective amount of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof.

In a further aspect the present invention provides the use of a compound of the  
20 Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, in the manufacture of a medicament for use in inhibiting TNF, IL-1, IL-6 or IL-8.

In a further aspect the present invention provides a method of inhibiting TNF, IL-1, IL-6 or IL-8 which comprises administering to a warm-blooded animal an effective amount of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester  
25 thereof.

In a further aspect the present invention provides the use of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, in the  
manufacture of a medicament for use in inhibiting TNF.

In a further aspect the present invention provides a method of inhibiting TNF which  
30 comprises administering to a warm-blooded animal an effective amount of a compound of the Formula I, or a pharmaceutically-acceptable salt or in vivo-cleavable ester thereof.

In a further aspect the present invention provides the use of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, in the manufacture of a medicament for use in the treatment of diseases or medical conditions mediated by p38 kinase.

5 In a further aspect the present invention provides a method of treating diseases or medical conditions mediated by p38 kinase which comprises administering to a warm-blooded animal an effective amount of a compound of the Formula I, or a pharmaceutically- acceptable salt or in-vivo-cleavable ester thereof.

---

In a further aspect the present invention provides the use of a compound of the  
10 Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, in the manufacture of a medicament for use in the production of a p38 kinase inhibitory effect.

In a further aspect the present invention provides a method of providing a p38 kinase inhibitory effect which comprises administering to a warm-blooded animal an effective amount of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-  
15 cleavable ester thereof.

In a further aspect the present invention provides the use of a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo- cleavable ester thereof, in the manufacture of a medicament for use in the treatment of rheumatoid arthritis, asthma, irritable bowel disease, multiple sclerosis, AIDS, septic shock, congestive heart failure, ischaemic  
20 heart disease or psoriasis.

In a further aspect the present invention provides a method of treating rheumatoid arthritis, asthma, irritable bowel disease, multiple sclerosis, AIDS, septic shock, congestive heart failure, ischaemic heart disease or psoriasis which comprises administering to a warm-blooded animal an effective amount of a compound of the Formula I, or a pharmaceutically-  
25 acceptable salt or in-vivo-cleavable ester thereof.

The compounds of this invention may be used in combination with other drugs and therapies used in the treatment of disease states which would benefit from the inhibition of cytokines, in particular TNF and IL-1. For example, the compounds of the Formula I could be used in combination with drugs and therapies used in the treatment of rheumatoid arthritis,  
30 asthma, irritable bowel disease, multiple sclerosis, AIDS, septic shock, congestive heart failure, ischaemic heart disease, psoriasis and the other disease states mentioned earlier in this specification.



For example, by virtue of their ability to inhibit cytokines, the compounds of the Formula I are of value in the treatment of certain inflammatory and non-inflammatory diseases which are currently treated with a cyclooxygenase-inhibitory non-steroidal anti-inflammatory drug (NSAID) such as indomethacin, ketorolac, acetylsalicylic acid, 5 ibuprofen, sulindac, tolmetin and piroxicam. Co-administration of a compound of the Formula I with a NSAID can result in a reduction of the quantity of the latter agent needed to produce a therapeutic effect. Thereby the likelihood of adverse side-effects from the NSAID such as gastrointestinal effects are reduced. Thus according to a further feature of the invention there is provided a pharmaceutical composition which comprises a compound of the 10 Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, in conjunction or admixture with a cyclooxygenase inhibitory non-steroidal anti-inflammatory agent, and a pharmaceutically-acceptable diluent or carrier.

The compounds of the invention may also be used with anti-inflammatory agents such as an inhibitor of the enzyme 5-lipoxygenase.

15 The compounds of the Formula I may also be used in the treatment of conditions such as rheumatoid arthritis in combination with antiarthritic agents such as gold, methotrexate, steroids and penicillinamine, and in conditions such as osteoarthritis in combination with steroids.

The compounds of the present invention may also be administered in degradative 20 diseases, for example osteoarthritis, with chondroprotective, anti-degradative and/or reparative agents such as Diacerhein, hyaluronic acid formulations such as Hyalan, Rumalon, Arteparon and glucosamine salts such as Antril.

The compounds of the Formula I may be used in the treatment of asthma in combination with antiasthmatic agents such as bronchodilators and leukotriene antagonists.

25 If formulated as a fixed dose such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically-active agent within its approved dosage range. Sequential use is contemplated when a combination formulation is inappropriate.

Although the compounds of the Formula I are primarily of value as therapeutic agents 30 for use in warm-blooded animals (including man), they are also useful whenever it is required to inhibit the effects of cytokines. Thus, they are useful as pharmacological standards for use in the development of new biological tests and in the search for new pharmacological agents.

The invention will now be illustrated in the following non-limiting Examples in which, unless otherwise stated:-

- (i) operations were carried out at ambient temperature, i.e. in the range 17 to 25°C and under an atmosphere of an inert gas such as argon unless otherwise stated;
- 5 (ii) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids by filtration;
- (iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck, Darmstadt,  
10 Germany or high pressure liquid chromatography (HPLC) was performed on C18 reverse phase silica, for example on a Dynamax C-18 60Å preparative reversed-phase column;
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) in general, the end-products of the Formula I have satisfactory microanalyses and  
15 their structures were confirmed by nuclear magnetic resonance (NMR) and/or mass spectral techniques; fast-atom bombardment (FAB) mass spectral data were obtained using a Platform spectrometer and, where appropriate, either positive ion data or negative ion data were collected; NMR chemical shift values were measured on the delta scale [proton magnetic resonance spectra were determined using a Varian Gemini 2000 spectrometer operating at a  
20 field strength of 300MHz or a Bruker AM250 spectrometer operating at a field strength of 250MHz]; the following abbreviations have been used: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad;
- (vi) intermediates were not generally fully characterised and purity was assessed by thin layer chromatographic, HPLC, infra-red (IR) and/or NMR analysis;
- 25 (vii) melting points are uncorrected and were determined using a Mettler SP62 automatic melting point apparatus or an oil-bath apparatus; melting points for the end-products of the Formula I were determined after crystallisation from a conventional organic solvent such as ethanol, methanol, acetone, ether or hexane, alone or in admixture; and
- 30 (viii) the following abbreviations have been used:-

DMF      N,N-dimethylformamide

DMSO dimethylsulphoxide

DMA N,N-dimethylacetamide.

---

**Example 1****6-Carbamoyl-2-chloro-4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]-pyrimidine**

A mixture of N-(3-amino-4-methylphenyl)-3-fluoro-5-morpholinobenzamide (0.33 g),  
 5 6-carbamoyl-2,4-dichloropyrimidine (0.212 g), N,N-di-isopropylethylamine (0.53 ml) and  
 n-butanol (5 ml) was stirred and heated to reflux for 36 hours. Water and isohexane were  
 added and the precipitated product was isolated and dried. There was thus obtained the title  
 compound (0.059 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.14 (s, 3H), 3.23 (m, 4H), 3.72 (m, 4H),  
 6.97 (d, 1H), 7.12 (d, 1H), 7.29 (m, 2H), 7.6 (d, 1H), 7.74 (s, 1H), 7.81 (s, 1H), 7.92 (s, 1H),  
 10 9.9 (s, 1H), 10.21 (s, 1H); Mass Spectrum: M-H<sup>+</sup> 483 & 485.

The N-(3-amino-4-methylphenyl)-3-fluoro-5-morpholinobenzamide used as a starting  
 material was obtained as follows :-

A solution of 3,5-difluorobenzoyl chloride (2.82 g) in methylene chloride (20 ml) was  
 added to a stirred mixture of 4-methyl-3-nitroaniline (2.28 g), triethylamine (4.35 ml) and  
 15 methylene chloride (80 ml). The resultant mixture was stirred at ambient temperature for  
 16 hours. The precipitate was isolated, washed with methylene chloride and dried. There was  
 thus obtained N-(4-methyl-3-nitrophenyl)-3,5-difluorobenzamide; NMR Spectrum:  
 (DMSO<sub>d</sub><sub>6</sub>) 2.43 (s, 3H), 7.43 (m, 2H), 7.63 (m, 2H), 7.95 (m, 2H), 8.43 (d, 1H), 10.42 (s,  
 1H); Mass Spectrum: M+H<sup>+</sup> 293.

20 A mixture of a portion (1 g) of the material so obtained and morpholine (5 ml) was  
 stirred and heated to 100°C for 48 hours and then to 120°C for 24 hours. The reaction  
 mixture was cooled and poured into water (100 ml). The resultant solid was isolated, washed  
 with water and dried. The material so obtained was purified by column chromatography on  
 silica using a 1:1 mixture of isohexane and ethyl acetate as eluent. There was thus obtained  
 25 N-(4-methyl-3-nitrophenyl)-3-fluoro-5-morpholinobenzamide as a solid (0.53 g); NMR  
Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.46 (s, 3H), 3.22 (t, 4H), 3.75 (t, 4H), 6.98 (m, 1H), 7.12 (d, 1H), 7.27  
 (s, 1H), 7.46 (d, 1H), 7.96 (m, 1H), 8.43 (d, 1H), 10.48 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 360.

A portion (0.483 g) of the compound so obtained was dissolved in ethyl acetate  
 (40 ml) and hydrogenated over 10% palladium-on-carbon catalyst (0.6 g) under an atmosphere  
 30 of hydrogen until the uptake of hydrogen ceased. The catalyst was removed by filtration and  
 the filtrate was evaporated. The residue was triturated under diethyl ether (25 ml). The  
 resultant solid was collected, washed with diethyl ether and dried. There was thus obtained


the required starting material (0.341 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.99 (s, 3H), 3.19 (t, 4H), 3.76 (t, 4H), 4.8 (s, 2H), 6.75 (d, 1H), 6.82 (d, 1H), 6.9 (d, 1H), 7.02 (s, 1H), 7.04 (d, 1H), 7.23 (s, 1H), 9.81 (s, 1H).

## 5 Example 2

Using an analogous procedure to that described in Example 1, the appropriate 4-chloropyrimidine was reacted with the appropriate aniline to give the compounds described in the following table. Unless otherwise stated, each aniline starting material is either commercially available or is readily prepared by standard methods from known materials.

10

Table I



Compound No.	(R <sup>1</sup> ) <sub>m</sub>	(R <sup>4</sup> ) <sub>p</sub>	Note
1	2-chloro	3-fluoro-5-morpholino	(a)
2	2-methylthio	3-fluoro-5-morpholino	(b)
3	6-chloro	3-fluoro-5-morpholino	(c)
4	6-chloro	3-morpholino	(d)
5	6-chloro	3-dimethylamino	(e)

### Notes

- (a) The reaction mixture was evaporated and the residue was purified by column chromatography on silica using increasingly polar mixtures of isohexane and ethyl acetate as eluent. The product so obtained gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.15 (s, 3H), 3.2 (m, 4H), 3.72 (m, 4H), 6.48 (d, 1H), 6.95 (m, 1H), 7.1 (m, 1H), 7.27 (m, 2H), 7.56 (m, 1H), 7.71 (m, 1H), 8.08 (d, 1H), 9.56 (s, 1H), 10.16 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 442.
- (b) The reaction mixture was heated to reflux for 24 hours. The resultant mixture was evaporated and the residue was purified by column chromatography on silica using a 19:1 mixture of methylene chloride and methanol as eluent. The product so obtained gave the

following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.16 (s, 3H), 2.37 (s, 3H), 3.21 (m, 4H), 3.74 (m, 4H), 6.29 (d, 1H), 6.96 (m, 1H), 7.1 (m, 1H), 7.25 (m, 2H), 7.5 (m, 1H), 7.82 (m, 1H), 8.03 (m, 1H), 9.04 (s, 1H), 10.13 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 454.

(c) The reaction mixture was evaporated and the residue was purified by column

5 chromatography on silica using increasingly polar mixtures of isohexane and ethyl acetate as eluent. The product so obtained gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.15 (s, 3H), 3.2 (m, 4H), 3.72 (m, 4H), 6.56 (s, 1H), 6.95 (m, 1H), 7.1 (m, 1H), 7.26 (m, 1H), 7.27 (s, 1H), 7.56 (m, 1H), 7.76 (m, 1H), 8.34 (s, 1H), 9.37 (s, 1H), 10.15 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 442.

10 (d) The reaction mixture was evaporated and the residue was purified by column chromatography on silica using increasingly polar mixtures of methylene chloride and methanol as eluent. The product so obtained gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.17 (s, 3H), 3.18 (m, 4H), 3.79 (m, 4H), 6.57 (s, 1H), 7.18 (m, 1H), 7.27 (m, 1H), 7.47 (m, 2H), 7.44 (m, 1H), 7.6 (m, 1H), 7.78 (m, 1H), 8.36 (s, 1H), 9.68 (s, 1H), 10.13 (s, 15 1H); Mass Spectrum: M+H<sup>+</sup> 424.

The N-(3-amino-4-methylphenyl)-3-morpholinobenzamide used as a starting material was prepared as follows :-

A mixture of ethyl 3-bromobenzoate (1.92 ml), morpholine (1.25 ml), 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.336 g), sodium tert-butoxide (1.615 g) and 20 tris(dibenzylideneacetone)dipalladium(0) (0.33 g) and toluene (25 ml) was stirred and heated to 90°C for 18 hours under argon. The reaction mixture was allowed to cool to ambient temperature and extracted with 1N aqueous hydrochloric acid. The aqueous phase was basified with concentrated sodium hydroxide solution and extracted with ethyl acetate. The organic phase was dried (MgSO<sub>4</sub>) and evaporated. The residual oil was 25 purified by column chromatography on silica gel using a 47:3 mixture of methylene chloride and methanol as eluent. There was thus obtained N-(3-morpholinobenzoyl)morpholine (0.45 g).

A mixture of the material so obtained, 5M sodium hydroxide solution (2.5 ml) and butanol (2 ml) was stirred and heated to 115°C for 18 hours. The mixture was evaporated 30 and the residue was acidified by the addition of 1N aqueous hydrochloric acid solution (12.5 ml). The resultant precipitate was isolated, washed with water and dried to give 3-morpholinobenzoic acid (0.15 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 3.1 (t, 4H), 3.73 (t, 4H),

7.19 (d, 1H), 7.32 (d, 1H), 7.38 (t, 1H), 7.42 (s, 1H).

Oxalyl chloride (0.14 ml) was added to a solution of 3-morpholinobenzoic acid (0.28 g) in methylene chloride (10 ml) which contained DMF (2 drops). The reaction mixture was stirred for 18 hours at ambient temperature. The mixture was evaporated and  
5 azeotroped with toluene to give 3-morpholinobenzoyl chloride (0.3 g); Mass Spectrum:  $M+H^+$  222.

A solution of 3-morpholinobenzoyl chloride (0.24 g) in methylene chloride (5 ml) was added to a stirred mixture of 4-methyl-3-nitroaniline (0.15 g), pyridine (0.24 ml) and methylene chloride (10 ml). The reaction mixture was stirred at ambient temperature for  
10 16 hours. The organic phase was washed with water and with a saturated aqueous sodium bicarbonate solution. The organic layer was dried ( $MgSO_4$ ) and evaporated. The residual solid was triturated under diethyl ether and the resultant solid was isolated and dried to give  
15 N-(3-nitro-4-methylphenyl)-3-morpholinobenzamide (0.28 g); NMR Spectrum: ( $DMSO-d_6$ ) 3.2 (t, 4H), 3.3 (s, 3H), 3.78 (t, 4H), 7.19 (s, 1H), 7.4 (m, 2H), 7.47 (d, 2H), 8.0 (d, 1H), 8.83 (s, 1H), 10.23 (s, 1H).

10% Palladium-on-carbon (0.035 g) was added to a stirred solution in methanol (40 ml) of the nitro compound so obtained (0.28 g) and the mixture was stirred at ambient temperature under 1 atmosphere pressure of hydrogen. After uptake of hydrogen had ceased, the catalyst was removed by filtration and the filtrate was evaporated to give  
20 N-(3-amino-4-methylphenyl)-3-morpholinobenzamide; NMR Spectrum: ( $DMSO-d_6$ ) 2.0 (s, 3H), 3.19 (t, 4H), 3.78 (t, 4H), 4.8 (s, 2H), 6.8 (q, 2H), 7.08 (s, 1H), 7.1 (d, 1H), 7.34 (m, 2H), 7.4 (s, 1H), 9.8 (s, 1H); Mass Spectrum:  $M+H^+$  312.

(e) The product gave the following data : NMR Spectrum: ( $DMSO-d_6$ ) 2.17 (s, 3H), 2.97 (s, 6H), 6.56 (m, 1H), 6.94 (m, 1H), 7.2-7.33 (m, 4H), 7.6 (m, 1H), 7.8 (m, 1H), 8.37 (s, 1H),  
25 9.39 (s, 1H), 10.1 (s, 1H); Mass Spectrum:  $M+H^+$  382 & 384.

The N-(3-amino-4-methylphenyl)-3-dimethylaminobenzamide used as starting material was prepared as follows :

Oxalyl chloride (13.0 ml) was added dropwise to a stirred mixture of 3-dimethylaminobenzoic acid (20.3 g) and DMF (a few drops) which had been cooled to  
30 0°C. The mixture was allowed to warm to ambient temperature and was stirred for 4 hours. The resultant mixture was evaporated and the residue was dissolved in methylene chloride (150 ml). 4-Methyl-3-nitroaniline (15.2 g) and triethylamine (27.9 ml) were

added in turn and the resultant mixture was stirred at ambient temperature for 16 hours. The reaction mixture was washed in turn with water, with a saturated aqueous solution of sodium bicarbonate and with a saturated aqueous sodium chloride solution, dried over magnesium sulphate and evaporated. The residue was triturated under a mixture of ethyl acetate and isohexane. The solid so obtained was filtered off and recrystallised from ethanol to give N-(3-nitro-4-methylphenyl)-3-dimethylamino-benzamide (6.1 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.46 (s, 3H), 2.95 (s, 6H), 6.92 (d, 1H), 7.22 (m, 2H), 7.32 (t, 1H), 7.45 (d, 1H), 7.97 (d, 1H), 8.53 (s, 1H), 10.43 (s, 1H).

After repetition of the previous reactions, a sample (8.25 g) was added to a stirred suspension of ammonium formate (17.4 g), and 10% palladium-on-carbon (1 g) in methanol (250 ml). The mixture was stirred and heated to reflux for 4 hours. The mixture was allowed to cool and then filtered. The filtrate was evaporated and water was added to the residue. The resultant solid was isolated and washed in turn with water, with ethyl acetate and with diethyl ether. The solid was dried in a vacuum oven at 40°C to give N-(3-amino-4-methylphenyl)-3-dimethylaminobenzamide (6.89 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.0 (s, 3H), 2.94 (s, 6H), 4.78 (s, 2H), 6.82 (m, 3H), 7.07 (s, 1H), 7.17 (m, 2H), 7.25 (m, 1H), 9.74 (s, 1H).

### Example 3

#### 20 **5-Amino-6-chloro-4-[5-(3-fluoro-5-morpholinobenzamido) 2-methylanilino]pyrimidine**

Ethereal hydrogen chloride solution (1M, 1.5 ml) was added to a mixture of 5-amino-4,6-dichloropyrimidine (0.512 g), N-(3-amino-4-methylphenyl)-3-fluoro-5-morpholinobenzamide (0.514 g) and n-butanol (10 ml) and the resultant mixture was stirred and heated to reflux for 24 hours. Di-isopropylamine (1 ml) was added and the mixture was allowed to cool to ambient temperature. Water and a mixture of diethyl ether and isohexane were added and the precipitated solid was collected and dried. There was thus obtained the title compound (0.373 g); NMR Spectrum: 2.12 (s, 3H), 3.22 (m, 4H), 3.74 (m, 4H), 5.28 (s, 2H), 6.87 (d, 1H), 7.11 (d, 1H), 7.22 (d, 1H), 7.27 (s, 1H), 7.52 (d, 1H), 7.69 (s, 1H), 7.73 (s, 1H), 8.03 (s, 1H), 10.15 (s, 1H); Mass Spectrum: M-H<sup>+</sup> 455 & 457.



**Example 4****4-[2-Methyl-5-(3-morpholinobenzamido)anilino]pyrimidine**

10% Palladium-on-carbon (0.02 g) was added to a mixture of 4-[2-methyl-5-(3-morpholinobenzamido)anilino]-6-chloropyrimidine (0.169 g), ammonium formate (0.165 g) and ethanol (5 ml) and the resultant mixture was stirred and heated to reflux for 18 hours. The catalyst was removed by filtration and the filtrate was evaporated. The residue was partitioned between methylene chloride and water and the organic phase was washed with brine, dried over sodium sulphate and evaporated. There was thus obtained the title compound (0.145 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.23 (s, 3H), 3.24 (m, 4H), 3.83 (m, 4H), 6.68 (m, 1H), 7.2 (m, 1H), 7.3 (m, 1H), 7.43 (m, 2H), 7.5 (m, 1H), 7.61 (m, 1H), 7.88 (m, 1H), 8.28 (m, 1H), 8.57 (s, 1H), 9.06 (s, 1H), 10.18 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 390.


**Example 5****4-[5-(3-Dimethylaminobenzamido)-2-methylanilino]-6-(3-pyrrolidin-1-ylpropylamino)pyrimidine**

A mixture of 4-[5-(3-dimethylaminobenzamido)-2-methylanilino]-6-chloropyrimidine (0.114 g), 3-pyrrolidin-1-ylpropylamine (0.95 ml), N,N-di-isopropylethylamine (0.063 ml) and n-butanol (2 ml) was stirred and heated to reflux for 18 hours. The mixture was evaporated and the residue was purified by column chromatography on silica using a 10:1 mixture of ethyl acetate and methanol as eluent. There was thus obtained the title compound (0.124 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.59 (m, 6H), 2.15 (s, 3H), 2.34 (m, 6H), 2.93 (s, 6H), 3.14 (m, 2H), 6.72 (m, 1H), 6.89 (m, 1H), 7.2 (m, 4H), 7.3 (m, 1H), 7.5 (m, 1H), 7.72 (m, 1H), 7.96 (m, 1H), 8.16 (s, 1H), 10.02 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 474.

**Example 6**

Using an analogous procedure to that described in Example 1, the appropriate 4-chloropyrimidine was reacted with the appropriate aniline to give the compounds described in the following table. Unless otherwise stated, each aniline starting material is either commercially available or is readily prepared by standard methods from known materials.

Table II



Compound No.	(R <sup>1</sup> ) <sub>m</sub>	R <sup>4</sup>	Note
1	2-chloro	morpholino	(a)
2	6-chloro	morpholino	(b)
3	5-bromo-2-methylthio	morpholino	(c)

### Notes

- 5 (a) The reaction mixture was evaporated and the residue was purified by column chromatography on silica using a 3:1 mixture of isohexane and ethyl acetate as eluent. The product so obtained gave the following data : NMR Spectrum: (DMSO-d<sub>6</sub>) 2.15 (s, 3H), 3.52 (m, 4H), 3.71 (m, 4H), 6.49 (m, 1H), 7.21 (s, 1H), 7.27 (m, 1H), 7.57 (m, 1H), 7.72 (m, 1H), 8.08 (m, 1H), 8.26 (m, 1H), 9.97 (s, 1H), 10.30 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 425 & 427.

- 10 The N-(3-amino-4-methylphenyl)-2-morpholinopyridine-4-carboxamide used as a starting material was prepared as follows :-

Triethylamine (31.8 ml) was added to a stirred mixture of 4-methyl-3-nitroaniline (15.8 g), 2-chloropyridine-4-carbonyl chloride (20 g) and methylene chloride (1 litre) and the resultant mixture was stirred at ambient temperature for 16 hours. The precipitate was isolated, washed with a saturated aqueous sodium bicarbonate solution and with methylene chloride and dried under vacuum at 40°C. There was thus obtained 2-chloro-N-(4-methyl-3-nitrophenyl)pyridine-4-carboxamide (10.2 g). The organic filtrate was washed with a saturated aqueous sodium bicarbonate solution, dried over magnesium sulphate and evaporated. The residue was triturated under methylene chloride and the resultant solid was isolated and dried under vacuum at 40°C. There was thus obtained a second crop (8.13 g) of 2-chloro-N-(4-methyl-3-nitrophenyl)pyridine-4-carboxamide; NMR Spectrum: (DMSO-d<sub>6</sub>) 2.48 (s, 3H), 7.51 (d, 1H), 7.86 (m, 1H), 7.96 (m, 2H), 8.49 (m, 1H), 8.64 (m, 1H), 10.85 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 292 and 294.

A mixture of the pyridine-4-carboxamide so produced and morpholine (250 ml) was stirred and heated to 100°C for 18 hours. The mixture was poured into water (250 ml) and stirred for 10 minutes. Methylene chloride (30 ml) was added and the resultant mixture was stirred for 30 minutes. The resultant solid was isolated, washed with

- 5 methylene chloride and dried in a vacuum oven at 40°C for 18 hours. There was thus obtained N-(4-methyl-3-nitrophenyl)-2-morpholinopyridine-4-carboxamide (17.34 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.48 (s, 3H), 3.52 (m, 4H), 3.71 (m, 4H), 7.1 (d, 1H), 7.25 (s, 1H), 7.49 (d, 1H) 7.97 (m, 1H), 8.29 (m, 1H), 8.49 (m, 1H), 10.62 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 343.

- 10 A mixture of a portion (8.5 g) of the material so obtained, 5% palladium-on-carbon catalyst (0.85 g) and methanol (600 ml) was stirred under an atmosphere pressure of hydrogen gas for 18 hours. Methylene chloride (400 ml) was added and the reaction mixture was filtered through diatomaceous earth. The filtrate was evaporated to give N-(3-amino-4-methylphenyl)-2-morpholinopyridine-4-carboxamide (6.41 g); NMR  
15 Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.01 (s, 3H), 3.52 (m, 4H), 3.73 (m, 4H), 4.83 (s, 2H), 6.78 (d, 1H), 6.84 (d, 1H) 7.04-7.08 (m, 2H), 7.2 (s, 1H), 8.24 (d, 1H), 9.95 (s, 1H); Mass Spectrum M+H<sup>+</sup> 313.

- (b) The reaction mixture was evaporated and the residue was purified by column chromatography on silica using increasingly polar mixtures of methylene chloride and  
20 methanol as eluent. The product so obtained gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.17 (s, 3H), 3.52 (m, 4H), 3.7 (m, 4H), 6.56 (s, 1H), 7.09 (m, 1H), 7.2 (m, 1H), 7.26 (m, 1H), 7.57 (m, 1H), 7.76 (m, 1H), 8.26 (m, 1H), 8.34 (s, 1H), 9.37 (s, 1H), 10.58 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 425.

- (c) The reaction mixture was heated to reflux for 18 hours. The mixture was cooled to  
25 ambient temperature and an excess of 1M ethereal hydrogen chloride solution was added. The resultant mixture was heated to reflux for 24 hours. After cooling, isohexane was added and the precipitated solid was collected by filtration. The product was obtained as a hydrochloride salt and gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.12 (s, 3H), 2.21 (s, 3H), 3.64 (m, 4H), 3.73 (m, 4H), 7.19 (d, 1H), 7.28 (d, 1H), 7.53 (s, 1H), 7.6 (m, 1H), 7.8 (s, 1H), 8.23  
30 (d, 1H), 8.35 (s, 1H), 8.92 (s, 1H), 10.57 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 515 & 517.

- (d) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) ; Mass Spectrum: M+H<sup>+</sup> .

**Example 7****4-[2-Methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine**

Using an analogous procedure to that described in Example 4, 6-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine was reacted with ammonium formate to give the title compound as a solid in 100% yield; NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.23 (s, 3H), 3.57 (m, 4H), 3.77 (m, 4H), 6.68 (m, 1H), 7.16 (m, 1H), 7.29 (m, 1H), 7.3 (m, 1H), 7.59 (m, 1H), 7.88 (m, 1H), 8.28 (m, 1H), 8.33 (m, 1H), 8.55 (s, 1H), 9.09 (s, 1H), 10.36 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 391.

---


10 **Example 8****4-[2-Methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-(N-methylpiperidin-4-yloxy)pyrimidine**

A mixture of 2-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.13 g), 4-hydroxy-N-methylpiperidine (0.087 g) and potassium tert-butoxide (1M in tert-butanol, 1.5 ml) was stirred and heated to 140°C for 1 hour. The mixture was partitioned between water and methylene chloride and the organic phase was washed with brine, dried over sodium sulphate and evaporated. There was thus obtained the title compound as a solid (0.11 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.45 (m, 2H), 1.88 (m, 2H), 1.97 (m, 2H), 2.05 (s, 3H), 2.15 (s, 3H), 2.23 (m, 2H), 3.5 (m, 4H), 5.7 (m, 4H), 4.77 (m, 1H), 6.28 (d, 1H), 6.89 (m, 1H), 7.1 (m, 1H), 7.21 (m, 1H), 7.24 (s, 1H), 7.47 (m, 1H), 7.9 (m, 1H), 7.98 (d, 1H), 8.25 (m, 1H), 8.9 (s, 1H), 10.24 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 504.

**Example 9**

25 Using an analogous procedure to that described in Example 8, the appropriate chloro-substituted pyrimidine was reacted with the appropriate alcohol to give the compounds described in the following table. Unless otherwise stated, each alcohol starting material is either commercially available or is readily prepared by standard methods from known materials.

Table III



No.	(R <sup>1</sup> ) <sub>m</sub>	R <sup>4</sup>	Note
1	2-(3-dimethylaminopropoxy)	morpholino	(a)
2	2-(2-dimethylaminopropoxy)	morpholino	(b)
3	2-(3-dimethylamino-2,2-dimethylpropoxy)	morpholino	(c)
4	2-[2-(N-methylpyrrolidin-2-yl)ethoxy]	morpholino	(d)
5	2-(N-isopropylpyrrolidin-3-yloxy)	morpholino	(e)
6	2-(N-tert-butoxycarbonylpiperidin-4-yloxy)	morpholino	(f)
7	2-(N-methylpyrrolidin-2-ylmethoxy)	morpholino	(g)
8	2-(N-methylpiperidin-3-ylmethoxy)	morpholino	(h)
9	2-(N,N'-dimethylpiperazin-2-ylmethoxy)	morpholino	(i)

### Notes

- 5 (a) The product gave the following data : NMR Spectrum: (DMSO-d<sub>6</sub>) 1.74 (m, 2H), 2.07 (s, 6H), 2.17 (s, 3H), 2.25 (t, 2H), 3.51 (m, 4H), 3.7 (m, 4H), 4.17 (t, 2H), 6.26 (d, 1H), 7.1 (m, 1H), 7.24 (s, 1H), 7.25 (m, 1H), 7.6 (m, 1H), 7.88 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 8.97 (s, 1H), 10.28 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 492.
- (b) The product gave the following data : NMR Spectrum: (DMSO-d<sub>6</sub>) 0.91 (d, 3H), 2.13 (s, 6H), 2.18 (s, 3H), 2.81 (m, 1H), 3.5 (m, 4H), 3.7 (m, 4H), 4.0 (m, 1H), 4.2 (m, 1H), 6.27 (d, 1H), 7.19 (m, 1H), 7.21 (s, 1H), 7.23 (m, 1H), 7.58 (m, 1H), 7.89 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 8.99 (s, 1H), 10.28 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 492.
- 10 (c) The product gave the following data : NMR Spectrum: (DMSO-d<sub>6</sub>) 0.88 (s, 6H), 2.15 (s, 6H), 1.18 (s, 3H), 3.34 (s, 2H), 3.52 (m, 4H), 3.71 (m, 4H), 3.94 (s, 2H), 6.23 (d, 1H), 7.1 (m, 1H), 7.22 (s, 1H), 7.24 (m, 1H), 7.49 (m, 1H), 7.91 (m, 1H), 8.0 (d, 1H), 8.28 (m, 1H), 9.01 (s, 1H), 10.29 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 520.
- 15

(d) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.37 (m, 1H), 1.52 (m, 1H), 1.79 (m, 1H), 1.91-2.28 (m, 4H), 2.13 (s, 3H), 2.18 (s, 3H), 2.7 (m, 1H), 2.86 (m, 1H), 3.5 (m, 4H), 3.7 (m, 4H), 4.17 (m, 1H), 6.29 (d, 1H), 7.1 (m, 1H), 7.21 (s, 1H), 7.23 (m, 1H), 7.49 (m, 1H), 7.91 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 8.98 (s, 1H), 10.27 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 518.

(e) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 0.93 (s, 3H), 0.97 (s, 3H), 1.8 (m, 1H), 2.05-2.36 (m, 3H), 2.17 (s, 3H), 2.55 (m, 1H), 2.65 (m, 1H), 2.75 (m, 1H), 3.5 (m, 4H), 3.7 (m, 4H), 5.08 (m, 1H), 6.3 (d, 1H), 7.1 (m, 1H), 7.21 (s, 1H), 7.23 (m, 1H), 7.5 (m, 1H), 7.85 (m, 1H), 7.99 (d, 1H), 8.27 (m, 1H), 8.97 (s, 1H), 10.28 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 518.

(f) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.47 (s, 9H), 1.58-1.8 (m, 4H), 1.98 (m, 2H), 2.04 (s, 3H), 3.06 (m, 2H), 3.58 (m, 4H), 3.81 (m, 4H), 5.15 (m, 1H), 6.25 (d, 1H), 6.68 (s, 1H), 6.9 (s, 1H), 7.11 (s, 1H), 7.26 (m, 2H), 7.83 (m, 1H), 7.92 (s, 1H), 8.07 (m, 1H), 8.3 (m, 1H); Mass Spectrum: M+H<sup>+</sup> 590.

15 The N-tert-butoxycarbonyl-4-hydroxypiperidine used as a starting material was obtained from a commercial source, for example from Neosystem, F67100, Strasbourg, France, or was prepared by the following procedure. A solution of di-tert-butyl dicarbonate (53.9 g) in methylene chloride (100 ml) was added dropwise to a stirred mixture of 4-hydroxypiperidine (25 g), triethylamine (50 ml) and methylene chloride (250 ml) which had  
20 been cooled to 0°C. The resultant mixture was allowed to warm to ambient temperature and was stirred for 18 hours. The mixture was evaporated and the residue was purified by chromatography on silica a 2:1 mixture of isohexane and ethyl acetate as eluent. The oil so obtained was dried under vacuum at 60°C to give N-tert-butoxycarbonyl-4-hydroxypiperidine as a white solid (49.1 g); NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.39 (s, 9H), 1.55 (m, 2H), 1.78 (m,  
25 2H), 2.95 (m, 2H), 3.76 (m, 2H).

(g) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.47-1.63 (m, 3H), 1.85 (m, 1H), 2.1 (m, 2H), 2.16 (s, 3H), 2.25 (s, 3H), 2.86 (m, 1H), 3.51 (m, 4H), 3.71 (m, 4H), 3.99 (m, 1H), 4.17 (m, 1H), 6.25 (d, 1H), 7.09 (m, 1H), 7.21 (s, 1H), 7.23 (m, 1H), 7.49 (m, 1H), 7.67 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 9.0 (s, 1H), 10.28 (s, 1H); Mass Spectrum:  
30 M+H<sup>+</sup> 504.

(h) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 0.92 (m, 1H), 1.4 (m, 1H), 1.6 (m, 3H), 1.79 (t, 1H), 1.9 (m, 1H), 2.08 (s, 3H), 2.18 (s, 3H), 2.69 (m, 1H), 3.35


(m, 1H), 3.52 (m, 4H), 3.7 (m, 4H), 4.03 (m, 2H), 6.27 (d, 1H), 7.10 (m, 1H), 7.20 (s, 1H), 7.22 (m, 1H), 7.47 (m, 1H), 7.92 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 8.99 (s, 1H), 10.28 (s, 1H); Mass Spectrum:  $M+H^+$  518.

(i) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.09 (t, 1H), 1.81 (t, 1H), 1.98 (t, 1H), 2.08 (s, 3H), 2.17 (s, 3H), 2.19 (s, 3H), 2.29 (m, 1H), 2.61 (m, 3H), 3.52 (m, 4H), 3.71 (m, 4H), 4.05 (m, 1H), 4.3 (m, 1H), 6.27 (d, 1H), 7.1 (m, 1H), 7.21 (s, 1H), 7.23 (m, 1H), 7.5 (m, 1H), 7.9 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 9.0 (s, 1H), 10.28 (s, 1H); Mass Spectrum:  $M+H^+$  533.

#### 10 Example 10

Using an analogous procedure to that described in Example 5, the appropriate chloro-substituted pyrimidine was reacted with the appropriate amine to give the compounds described in the following table. Unless otherwise stated, each amine starting material is either commercially available or is readily prepared by standard methods from known materials.

**Table IV**



No.	(R <sup>1</sup> ) <sub>m</sub>	R <sup>4</sup>	Note
1	2-morpholino	morpholino	(a)
2	2-(2,6-dimethylmorpholin-4-yl)	morpholino	(b)
3	2-(tetrahydro-4H-1,4-thiazin-4-yl)	morpholino	(c)
4	2-(4-methylpiperazin-1-yl)	morpholino	(d)
5	2-(4-isopropylpiperazin-1-yl)	morpholino	(e)
6	2-(4- <u>tert</u> -butoxycarbonylpiperazin-1-yl)	morpholino	(f)
7	2-(4-hydroxypiperidin-1-yl)	morpholino	(g)
8	2-[N-(N-methylpiperidin-4-yl)amino]	morpholino	(h)

9	2-[N-methyl-N-(N-methylpiperidin-4-yl)amino]	morpholino	(i)
10	2-[N-(N-ethylpiperidin-3-yl)amino]	morpholino	(j)
11	2-[N-(N-ethylpyrrolidin-2-ylmethyl)amino]	morpholino	(k)
12	2-[2-(2-ethylimidazol-1-yl)ethylamino]	morpholino	(l)
13	2-[2-(2-oxoimidazolidin-1-yl)ethylamino]	morpholino	(m)
14	2-(2-imidazol-1-ylethylamino)	morpholino	(n)
15	2-[3-(1,2,4-triazol-1-yl)propylamino]	morpholino	(o)
16	2-(3-pyrrolidin-1-ylpropylamino)	morpholino	(p)
17	2-[3-(2-oxopyrrolidin-1-yl)propylamino]	morpholino	(q)
18	2-[2-(N-methylpyrrolidin-2-yl)ethylamino]	morpholino	(r)
19	2-(3-imidazol-1-ylpropylamino)	morpholino	(s)
20	2-(3-morpholinopropylamino)	morpholino	(t)
21	2-(3-diethylaminopropylamino)	morpholino	(u)
22	2-(3-dimethylamino-2,2-dimethylpropylamino)	morpholino	(v)
23	2-(3-amino-2-hydroxypropylamino)	morpholino	(w)
24	2-(4-dimethylaminobutylamino)	morpholino	(x)
25	2-(4-imidazol-1-ylbutylamino)	morpholino	(y)
26	6-[N-methyl-N-(N-methylpiperidin-4-yl)amino]	morpholino	(z)
27	6-[N-(N-ethylpiperidin-3-yl)amino]	morpholino	(aa)
28	6-(3-imidazol-1-ylpropylamino)	morpholino	(bb)
29	6-[3-(4-methylpiperazin-1-yl)propylamino]	morpholino	(cc)
30	6-[N-methyl-N-(3-morpholinopropyl)amino]	morpholino	(dd)
31	6-(3-pyrrolidin-1-ylpropylamino)	morpholino	(ee)
32	6-(4-dimethylaminobutylamino)	morpholino	(ff)
33	6-(3-dimethylamino-2,2-dimethylpropylamino)	morpholino	(gg)
34	6-(3-diethylaminopropylamino)	morpholino	(hh)
35	6-(3-dimethylaminopropylamino)	morpholino	(ii)
36	6-[3-(2-oxopyrrolidin-1-yl)propylamino]	morpholino	(jj)

### Notes

(a) The product gave the following data : Mass Spectrum:  $M+H^+$  476.



- (b) The product gave the following data : Mass Spectrum:  $M+H^+$  504.
- (c) The product gave the following data : Mass Spectrum:  $M+H^+$  492.
- (d) The product gave the following data : NMR Spectrum: (DMSO $d_6$ ) 2.13 (s, 3H), 2.16 (s, 3H), 2.27 (m, 4H), 3.52 (m, 4H), 3.62 (m, 4H), 3.72 (m, 4H), 5.92 (m, 1H), 7.08 (m, 1H),  
5 7.18 (m, 1H), 7.2 (m, 1H), 7.36 (m, 1H), 7.87 (m, 1H), 8.02 (m, 1H), 8.27 (m, 1H), 8.47 (s, 1H) 10.19 (s, 1H); Mass Spectrum:  $M+H^+$  489.
- (e) The product gave the following data : Mass Spectrum:  $M+H^+$  517.
- 
- (f) The product gave the following data : Mass Spectrum:  $M+H^+$  575.
- (g) The product gave the following data : Mass Spectrum:  $M+H^+$  490.
- 10 (h) The product gave the following data : NMR Spectrum: (DMSO $d_6$ ) 1.29 (m, 2H), 1.93 (m, 2H), 2.27 (s, 3H), 2.57 (m, 3H), 2.6 (m, 1H), 2.94 (t, 2H), 3.58 (m, 4H), 5.88 (m, 4H), 4.64 (m, 2H), 5.88 (m, 1H), 6.28 (m, 1H), 6.89 (m, 1H), 7.1 (s, 1H), 7.21 (m, 1H), 7.27 (s, 1H), 7.33 (m, 1H), 7.8 (s, 1H), 7.93 (m, 1H), 7.98 (m, 1H), 8.29 (m, 1H); Mass Spectrum:  $M+H^+$  503.
- 15 (i) The product gave the following data : NMR Spectrum: (DMSO $d_6$ ) 1.68 (m, 2H), 1.88 (m, 2H), 2.09 (m, 2H), 2.27 (s, 3H), 2.28 (s, 3H), 2.91 (m, 2H), 3.03 (s, 3H), 3.6 (m, 4H), 3.82 (m, 4H), 4.6 (m, 1H), 5.89 (d, 1H), 6.25 (s, 1H), 6.9 (d, 1H), 7.1 (s, 1H), 7.24 (m, 1H), 7.32 (m, 1H), 7.85 (m, 1H), 8.0 (d, 1H), 8.3 (d, 1H); Mass Spectrum:  $M+H^+$  517.
- (j) The product gave the following data : NMR Spectrum: (DMSO $d_6$ ) 0.96 (t, 3H), 1.4  
20 (m, 2H), 1.6 (m, 2H), 1.88 (m, 2H), 2.28 (s, 3H), 2.3 (m, 4H), 3.58 (m, 4H), 5.82 (m, 4H), 4.03 (m, 1H), 5.11 (m, 1H), 5.92 (m, 1H), 6.3 (s, 1H), 6.94 (m, 1H), 7.1 (s, 1H), 7.2 (m, 1H), 7.28 (s, 1H), 7.6 (m, 1H), 7.93 (m, 1H), 8.28 (m, 1H), 8.54 (m, 1H); Mass Spectrum:  $M+H^+$  517.
- (k) The product gave the following data : Mass Spectrum:  $M+H^+$  517.
- 25 (l) The 2-(2-ethylimidazol-1-yl)ethylamine, used as a starting material, is disclosed in Chemical Abstracts, volume 108, abstract 151535. The product gave the following data : Mass Spectrum:  $M+H^+$  528.
- (m) The product gave the following data : Mass Spectrum:  $M+H^+$  518.
- (n) The 2-imidazol-1-ylethylamine, used as a starting material, is disclosed in J. Medicinal  
30 Chemistry, 1986, 29, 523-530. The product gave the following data : Mass Spectrum:  $M+H^+$  500.

- (o) The 3-(1,2,4-triazol-1-yl)propylamine, used as a starting material, is disclosed in J. Medicinal Chemistry, 1986, 29, 523-530. The product gave the following data : Mass Spectrum:  $M+H^+$  515.
- (p) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.59 (m, 6H), 2.18 (s, 3H), 2.3 (m, 6H), 3.19 (m, 2H), 3.51 (m, 4H), 3.71 (m, 4H), 5.87 (m, 1H), 7.09 (m, 1H), 7.18 (m, 1H), 7.22 (m, 1H), 7.43 (m, 1H), 7.77 (m, 1H), 7.92 (m, 1H), 8.26 (m, 1H), 8.31 (s, 1H), 10.17 (s, 1H); Mass Spectrum:  $M+H^+$  517.
- 
- (q) The product gave the following data : Mass Spectrum:  $M+H^+$  531.
- (r) The product gave the following data : Mass Spectrum:  $M+H^+$  517.
- 10 (s) The product gave the following data : Mass Spectrum:  $M+H^+$  514.
- (t) The product gave the following data : Mass Spectrum:  $M+H^+$  533.
- (u) The product gave the following data : Mass Spectrum:  $M+H^+$  519.
- (v) The product gave the following data : Mass Spectrum:  $M+H^+$  519.
- (w) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.57 (s, 3H), 2.28 (m, 2H), 2.23 (m, 2H), 3.52 (m, 4H), 3.72 (m, 4H), 5.86 (m, 1H), 6.32 (m, 1H), 7.08 (m, 1H), 7.18 (m, 1H), 7.21 (m, 1H), 7.47 (m, 1H), 7.79 (m, 1H), 7.85 (m, 1H), 8.27 (m, 1H), 8.41 (s, 1H), 10.17 (s, 1H); Mass Spectrum:  $M+H^+$  479.
- (x) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 2.57 (s, 3H), 2.28 (m, 2H), 2.23 (m, 2H), 3.52 (m, 4H), 3.72 (m, 4H), 5.86 (m, 1H), 6.32 (m, 1H), 7.08 (m, 1H), 7.18 (m, 1H), 7.21 (m, 1H), 7.47 (m, 1H), 7.79 (m, 1H), 7.85 (m, 1H), 8.27 (m, 1H), 8.41 (s, 1H), 10.17 (s, 1H); Mass Spectrum:  $M+H^+$  505.
- 20 (y) The 4-imidazol-1-ylbutylamine, used as a starting material, is disclosed in J. Medicinal Chemistry, 1987, 30, 185-193. The product gave the following data : Mass Spectrum:  $M+H^+$  528.
- (z) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.66 (m, 2H), 2.86 (m, 4H), 2.12 (m, 2H), 2.26 (s, 3H), 2.28 (s, 3H), 2.83 (s, 3H), 2.9 (m, 2H), 3.57 (m, 4H), 3.82 (m, 4H), 5.79 (s, 1H), 6.38 (s, 1H), 6.89 (m, 1H), 7.07 (m, 1H), 7.14 (m, 1H), 7.22 (m, 1H), 7.92 (m, 1H), 8.03 (s, 1H), 8.3 (m, 1H); Mass Spectrum:  $M+H^+$  517.
- (aa) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.06 (m, 3H), 1.52 (m, 2H), 1.72 (m, 4H), 2.26 (s, 3H), 2.28 (m, 6H), 2.59 (m, 1H), 3.58 (m, 4H), 3.83 (m, 4H), 5.58 (s, 1H), 6.3 (s, 1H), 6.92 (m, 1H), 7.1 (m, 1H), 7.22 (m, 1H), 7.33 (m, 1H), 7.78 (m, 1H), 7.88 (s, 1H), 8.18 (s, 1H), 8.31 (m, 1H); Mass Spectrum:  $M+H^+$  517.
- 30

(bb) The product gave the following data : Mass Spectrum:  $M+H^+$  514.

(cc) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.61 (m, 2H), 2.12 (s, 3H), 2.17 (s, 3H), 2.28 (m, 5H), 3.13 (m, 2H), 3.52 (m, 4H), 3.72 (m, 4H), 5.46 (s, 1H), 6.78 (m, 1H), 7.1 (m, 1H), 7.19 (m, 1H), 7.23 (m, 1H), 7.5 (m, 1H), 7.74 (m, 1H), 7.98 (m, 1H), 8.2 (m, 1H), 8.27 (m, 1H), 10.25 (s, 1H); Mass Spectrum:  $M+H^+$  546.

(dd) The product gave the following data : Mass Spectrum:  $M+H^+$  547.

The N-(3-methylaminopropyl)morpholine, used as a starting material, was obtained by bubbling methylamine gas for 10 minutes through a solution of 3-morpholinopropyl chloride (3.27 g) in ethanol (30 ml). The resultant mixture was allowed to stand for 1 hour and then evaporated. The residual oil was triturated under diethyl ether and the resultant solid was dried under vacuum. There was thus obtained the required starting material (1.58 g).

(ee) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.62 (m, 6H), 2.15 (s, 3H), 2.36 (m, 6H), 3.14 (m, 2H), 3.52 (m, 4H), 3.72 (m, 4H), 6.72 (m, 1H), 7.08 (m, 1H), 7.2 (m, 1H), 7.22 (s, 1H), 7.48 (m, 1H), 7.76 (m, 1H), 7.96 (s, 1H), 8.17 (s, 1H), 8.26 (m, 1H); Mass Spectrum:  $M+H^+$  517.

(ff) The product gave the following data : Mass Spectrum:  $M+H^+$  505.

(gg) The product gave the following data : Mass Spectrum:  $M+H^+$  519.

(hh) The product gave the following data : Mass Spectrum:  $M+H^+$  519.

(ii) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.56 (m, 2H), 2.09 (s, 6H), 2.17 (s, 3H), 3.14 (m, 2H), 3.5 (m, 4H), 3.7 (m, 4H), 5.46 (s, 1H), 6.75 (m, 1H), 7.08 (m, 1H), 7.2 (m, 1H), 7.23 (m, 1H), 7.5 (m, 1H), 7.75 (m, 1H), 7.98 (m, 1H), 8.2 (m, 1H), 8.24 (m, 1H), 10.24 (s, 1H); Mass Spectrum:  $M+H^+$  491.

(jj) The product gave the following data : NMR Spectrum: (DMSO<sub>d</sub><sub>6</sub>) 1.63 (t, 2H), 1.87 (t, 2H), 2.16 (s, 3H), 3.19 (t, 4H), 3.3 (t, 4H), 3.51 (m, 4H), 3.7 (m, 4H), 5.47 (s, 1H), 6.74 (m, 1H), 7.1 (m, 1H), 7.2 (m, 1H), 7.22 (s, 1H), 7.5 (m, 1H), 7.73 (m, 1H), 7.98 (s, 1H), 8.26 (m, 2H), 10.24 (s, 1H); Mass Spectrum:  $M+H^+$  531.

### Example 11

4-[2-Methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-  
2-(piperidin-4-yloxy)pyrimidine

Trifluoroacetic acid (5 ml) was added to a solution of  
2-(1-tert-butoxycarbonylpiperidin-4-yloxy)-4-[2-methyl-5-(2-morpholinopyrid-

4-ylcarbonylamino)anilino]pyrimidine (0.212 g) in methylene chloride (5 ml) and the resultant mixture was stirred at ambient temperature for 2 hours. The mixture was evaporated and the residue was partitioned between methylene chloride and water. The aqueous phase was passed through an ion exchange column (isolute SCX column from International Sorbent Technology Limited, Hengoed, Mid-Glamorgan, UK) using a 99:1 mixture of methanol and a saturated aqueous ammonium hydroxide solution as eluent. The product so obtained was triturated under diethyl ether and there was thus obtained the title compound as a solid

(0.065 g); NMR Spectrum: (DMSO-d<sub>6</sub>) 1.2 (m, 2H), 1.87 (m, 2H), 2.17 (s, 3H), 2.19 (s, 3H), 2.27 (m, 2H), 2.95 (m, 2H), 3.5 (m, 4H), 3.7 (m, 4H), 4.85 (m, 1H), 6.27 (d, 1H), 7.1 (m, 1H), 7.21 (s, 1H), 7.23 (m, 1H), 7.5 (m, 1H), 7.9 (m, 1H), 8.0 (d, 1H), 8.27 (m, 1H), 9.0 (s, 1H), 10.28 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 490.

### Example 12

4-[2-Methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-(piperazin-1-yloxy)pyrimidine trihydrochloride

A mixture of 2-(4-tert-butoxycarbonylpiperazin-1-yl)-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.184 g) and a saturated solution of hydrogen chloride in ethyl acetate (20 ml) was stirred at ambient temperature for 2 hours. The resultant precipitate was collected and washed with diethyl ether. There was thus obtained the title compound (0.248 g); NMR Spectrum: (DMSO-d<sub>6</sub>) 2.22 (s, 3H), 3.2 (m, 4H), 3.65 (m, 4H), 3.72 (m, 4H), 3.95 (m, 4H), 7.2 (d, 1H), 7.28 (d, 1H), 7.49 (m, 1H), 7.58 (s, 1H), 7.9 (d, 1H), 8.14 (m, 1H), 8.2 (d, 1H), 9.5 (m, 1H), 10.5 (s, 1H), 10.75 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 475.

### 25 Example 13

2-Allylamino-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine

A mixture of 2-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.2 g) and allylamine (0.4 ml) was stirred and heated to reflux for 18 hours. The mixture was evaporated and the residue was purified by column chromatography on silica using a 9:1 mixture of methylene chloride and methanol as eluent. There was thus obtained the title compound as a solid (0.135 g); NMR Spectrum: (DMSO-d<sub>6</sub>) 2.17 (s, 3H), 3.5 (m, 4H), 3.7 (m, 4H), 3.82 (m, 2H), 4.95 (m, 1H), 5.05 (m, 1H), 5.85 (m,

2H), 6.57 (br m, 1H), 7.07 (m, 1H), 7.17 (m, 2H), 7.43 (m, 1H), 7.78 (m, 1H), 7.87 (br m, 1H), 8.24 (m, 1H), 8.37 (s, 1H), 10.18 (s, 1H); Mass Spectrum:  $M+H^+$  446.

#### Example 14

##### 5 2-(N-Allyl-N-methylamino)-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine

A mixture of 2-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.2 g) and N-allyl-N-methylamine (0.25 ml) was stirred and heated to 75°C for 3 hours. The mixture was cooled to ambient temperature and  
 10 purified by column chromatography on silica using a 9:1 mixture of methylene chloride and methanol as eluent. The resultant oil was triturated under a mixture of isohexane and diethyl ether. There was thus obtained the title compound as a solid (0.055 g); NMR Spectrum: (DMSO- $d_6$ ) 2.17 (s, 3H), 2.95 (s, 3H), 3.5 (m, 4H), 3.7 (m, 4H), 4.1 (d, 2H), 5.04 (m, 2H), 5.75 (m, 1H), 5.91 (m, 1H), 7.07 (m, 1H), 7.17 (m, 2H), 7.19 (m, 1H), 7.86 (m, 1H), 7.98 (m, 1H),  
 15 8.25 (m, 1H), 8.42 (s, 1H), 10.17 (s, 1H); Mass Spectrum:  $M+H^+$  460.

#### Example 15

##### 2-Isopropylamino-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine

20 A mixture of 2-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.2 g) and isopropylamine (5 ml) was heated to 100°C in a sealed tube for 24 hours. The mixture was evaporated and the residue was partitioned between methylene chloride and water. The organic phase was washed with brine, dried over sodium sulphate and evaporated. The resultant solid was recrystallised from tert-butyl methyl  
 25 ether to give the title compound (0.145 g); NMR Spectrum: 1.04 (s, 3H), 1.06 (s, 3H), 2.17 (s, 3H), 3.5 (m, 4H), 3.7 (m, 4H), 3.97 (m, 1H), 5.82 (m, 1H), 6.2 (br m, 1H), 7.07 (m, 1H), 7.19 (m, 2H), 7.45 (m, 1H), 7.78 (m, 1H), 7.87 (br m, 1H), 8.25 (m, 1H), 8.31 (s, 1H), 10.18 (s, 1H); Mass Spectrum:  $M+H^+$  448.

**Example 16****2-Amino-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine**

A mixture of 2-allylamino-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.151 g), methanesulphonic acid (0.1 ml) and 30% palladium-on-carbon (0.034 g) in DMA (1 ml) was stirred and heated to 140°C for 24 hours. The mixture was cooled to ambient temperature and filtered. A mixture of methylene chloride and isohexane was added to the filtrate. The resultant precipitate was isolated and purified by column chromatography on silica using a 89:10:1 mixture of methylene chloride, methanol and a saturated aqueous ammonium hydroxide solution as eluent. There was thus obtained the title compound as a solid (0.01 g); NMR Spectrum: 2.16 (s, 3H), 3.47 (m, 4H), 3.7 (m, 4H), 4.78 (br m, 4H), 5.98 (m, 1H), 7.01 (m, 1H), 7.16 (m, 2H), 7.39 (m, 1H), 7.65 (m, 1H), 7.72 (m, 1H), 8.17 (m, 1H); Mass Spectrum: M+H<sup>+</sup> 406.

**Example 17****2-(3-Dimethylamino-2-hydroxypropylamino)-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine**

Formaldehyde (40% solution in water, 0.2 ml) was added to a suspension of 2-(3-amino-2-hydroxypropylamino)-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine (0.1 g) and sodium cyanoborohydride (0.04 g) in a 3:1 mixture of ethanol and water (16 ml). Acetic acid (0.5 ml) was added to take the pH to approximately 4. The mixture was stirred at ambient temperature for 3 hours. The ethanol was evaporated and a saturated aqueous sodium bicarbonate solution was added to the residue to bring the pH to 8. The mixture was partitioned between methylene chloride and water. The organic phase was washed with brine, dried over sodium sulphate and evaporated to give the title compound as a solid (0.073 g); NMR Spectrum: 2.11 (s, 3H), 2.16 (s, 3H), 3.11 (m, 2H), 3.52 (m, 4H), 3.7 (m, 4H), 5.86 (m, 1H), 6.24 (m, 1H), 7.08 (m, 1H), 7.18 (m, 1H), 7.22 (m, 1H), 7.47 (m, 1H), 7.78 (m, 1H), 7.88 (m, 1H), 8.26 (m, 1H), 8.21 (s, 1H), 10.19 (s, 1H); Mass Spectrum: M+H<sup>+</sup> 507.

**This Page Blank (uspto)**

---

PCT/GB08/03929  
10/10/00  
AobraZeneca

---

**This Page Blank (uspto)**